arXiv:1202.2269v1 [math.KT] 10 Feb 2012

On the conjectural Leibniz cohomology for groups

Simon Covez

Abstract

The goal of this paper is to present results which are consistent with conjectures about the *Leibniz (co)homology for discrete groups*, enounced by J. L. Loday in [Lod03]. We show that the *rack cohomology* has properties very closed from those which are expected for the conjectural Leibniz cohomology. In particular we define a graded *dendriform* algebra structure on the rack cohomology, and we construct a graded associative algebra morphism from the group cohomology to the rack cohomology.

Introduction

Problems and results. The starting point of this article are conjectures given by J.-L. Loday in [Lod03]. In this article J.-L. Loday conjectures the existence of a Leibniz homology $HL_\bullet(-)$ for groups and some properties satisfied by it. In this paper we focus on two properties

1. $HL_\bullet(G)$ is a graded Zinbiel coalgebra (and, *a fortiori*, a graded commutative coalgebra),
2. There exists a morphism of graded cocommutative algebra from $HL_\bullet(G)$ to the usual group homology $H_\bullet(G)$.

Here we work in the cohomological context (because it is easier to define a product than a coproduct) and we show two results which are consistent with these conjectures $\big($Theorem (4.5) and Theorem (5.8)$\big)$. We prove that there exists a cohomology theory $HR^\bullet$, defined for groups, and satisfying :

1. $HR^\bullet(G)$ is a graded dendriform algebra (and, *a fortiori*, a graded associative algebra),
2. There exists a morphism of graded associative algebra from $H^\bullet(G)$ to $HR^\bullet(G)$.

To understand what is $HR^\bullet(-)$ and where it comes from, it is good to know the reason of these conjectures.

Algebraic K-theory and additive K-theory. The starting point is a program of J.-L. Loday exposed in [Lod03] which intend to solve two problems arising in algebraic K-theory using the knowledge of the solutions in the additive K-theory context. These two problems are :

1. Have a small presentation of all the algebraic K-groups of a field, or, at least, a small presentation of a chain complex whose homology would be these K-groups. (By small, it essentially means no matrices in the presentation),
2. Determine the obstructions to the periodicity.

Rationally, the algebraic K-theory $K_\bullet(A)_{\mathbb{Q}}$ of a ring with unit A can be defined using the group homology of $GL(A)$, as the primitive part of the graded connected cocommutative Hopf algebra $H_\bullet(GL(A),\mathbb{Q})$, hence, using the Milnor-Moore theorem, we have the following graded isomorphism

$$H_\bullet(GL(A),\mathbb{Q}) \simeq \Lambda(K_\bullet(A)_{\mathbb{Q}}).$$

Using this definition, the additive K-theory can be seen as the "tangent" version of algebraic K-theory. Let $\mathbb{K}$ be a field of characteristic zero and A be a unital $\mathbb{K}$-algebra. The additive K-theory $K^+_\bullet(A)$ is defined as the algebraic K-theory, replacing the group $GL(A)$ by its "tangent" Lie algebra $\mathfrak{gl}(A)$, and the group homology by the Lie algebra homology. Therefore $K^+_\bullet(A)$ is defined as the primitive part of the graded connected cocommutative Hopf algebra $H_\bullet(\mathfrak{gl}(A),\mathbb{K})$, and by the Milnor-Moore theorem we have the following graded isomorphism

$$H_\bullet(\mathfrak{gl}(A),\mathbb{K}) \simeq \Lambda\big(K^+_\bullet(A)\big).$$

In the additive K-theory context the solutions of our problems are given by :

1. The *cyclic homology* $HC_\bullet$ by the *Loday-Quillen-Tsygan Theorem* (cf.[LQ81, FT87])

   Theorem 0.1 (Loday-Quillen-Tsygan)**.** *Let $\mathbb{K}$ be a field containing $\mathbb{Q}$ and A be a unital associative $\mathbb{K}$-algebra. Then there is a natural isomorphism :*

   $$K^+_n(A) \simeq HC_{n-1}(A),\ n \geq 1.$$

2. The *Hoschchild homology* $HH_\bullet$ by the *Connes' periodicity exact sequence*

   $$\cdots \to HH_n(A) \to HC_n(A) \to HC_{n-2}(A) \to HH_{n-1}(A) \to \cdots .$$

The following theorem is the starting point of this program. In the same way as the Loday-Quillen-Tsygan Theorem links $HC_\bullet(A)$ and $\mathfrak{gl}(A)$, the *Loday-Cuvier Theorem* (cf. [Lod98, Cuv]) gives a link between $HH_\bullet(A)$ and $\mathfrak{gl}(A)$.

Theorem 0.2 (Loday-Cuvier)**.** *For any associative unital algebra A over a characteristic zero field $\mathbb{K}$ there is an isomorphism of graded modules :*

$$HL_\bullet(\mathfrak{gl}(A),\mathbb{K}) \simeq T\big(HH_{\bullet-1}(A)\big).$$

In this theorem $HL_\bullet(-)$ is the Leibniz homology, the homology theory naturally associated to the *Leibniz algebras* (cf [Lod93, Lod97, Lod98]). The Leibniz algebras are a non-commutative version of Lie algebras and their homology gives new invariants for Lie algebras. The existence of $HL^\bullet(-)$ and its following properties (cf. [Lod95]) can explain the reason of the conjectures :

1. $HL^\bullet(\mathfrak{g})$ is a graded Zinbiel algebra,
2. There exists a morphism of graded commutative algebra from $H^\bullet(\mathfrak{g})$ to $HL^\bullet(\mathfrak{g})$.

The Lie algebra homology being the "tangent" version of group homology, the Leibniz algebra homology should be the "tangent" version of the expected homology $HR_\bullet(-)$. This hypothesis leads J.-L. Loday to the formulation of the *coquecigrue problem* (cf. [Lod93, Lod03, Lod12]).

The coquecigrue problem. There are two ways to construct a Lie algebra from a group. One is to consider the graded abelian group $\oplus_n G^{(n)}/G^{(n+1)}$ associated to the descending central series $\{G^{(n)} = [G, G^{(n-1)}]\}_{n\in\mathbb{N}}$. This object is provided with a Lie algebra structure where the bracket is induced by the commutator in G, the Jacobi identity being a consequence of the so-called *Phillip Hall relation.*

Another way is to consider the tangent space at the neutral element of a Lie group. The bracket being induced by the conjugation in the group, and the Leibniz identity by the autodistributivity of the conjugation, that is, the relation :

$$x \rhd (y \rhd z) = (x \rhd y) \rhd (x \rhd z) \text{ where } x \rhd y = xyx^{-1}.$$

The coquecigrue problem is the research of the generalization of one of these constructions for Leibniz algebras. Therefore the problem is to know if there exists a mathematical object, dubbed coquecigrue, generalizing groups, such that a Leibniz algebras is naturally associated to it (by one of these constructions) ?

In [Cov11] a local solution to the coquecigrue problem has been done for the second construction using *racks*, especially the links between (Lie) rack cohomology $HR^\bullet(-)$ and Leibniz algebra cohomology $HL^\bullet(-)$. Originally, racks were defined by knot theorists to create invariants for knots (cf. [CS] and references therein), and it is M. K. Kinyon who gives the first in [Kin07] the relation between Lie racks and Leibniz algebras.

A group being a rack it is natural to suppose that the rack cohomology is our conjectural Leibniz cohomology $HR^\bullet(-)$, and so, to study the existence of an algebraic structure on it and its link with the usual group cohomology.

The plan for this article is the following :

Section 1 : Dendriform and Zinbiel algebras. This section is based on [Lod01]. We recall the definitions of dendriform and Zinbiel algebras and give some examples. We recall also the relations between these algebras and others types of algebras such as *associative, commutative, Lie, Leibniz and diassociative* algebras.

Section 2 : Racks. Racks are sets provided with a product which encapsuled some properties of the conjugation in a group (especially the *distributivity*).This section is based on [FR] for the general theory of racks and on [EG03] for the definition of the rack cohomology $HR^\bullet(-)$. We recall some basic definitions about racks and their cohomology.

Section 3 : The category of trunks. *Trunks* have been introduced by R. Fenn, C. Rourke and B. Sanderson in [FRS95] in order to define a classifying space for racks. Trunks are objects loosely analogeous to category. A category can be seen as a graph with a fixed set of triangle (the graph of the composition) satisfying the associativity condition.

$$\begin{tikzcd}
 & C \\
A \arrow[ur, "x_1\cdot x_2"] \arrow[r, "x_1"'] & B \arrow[u, "x_2"']
\end{tikzcd}
\qquad
\begin{tikzcd}
 & & D \\
 & & C \arrow[u, "x_3"'] \\
A \arrow[uurr, "x_1\cdot(x_2\cdot x_3)=(x_1\cdot x_2)\cdot x_3"] \arrow[urr, dotted, "x_1\cdot x_2"] \arrow[r, "x_1"'] & B \arrow[uur, "x_2\cdot x_3"] \arrow[ur, "x_2"'] &
\end{tikzcd}$$

Like a category, a trunk is a graph with a fixed set of *cubes*

$$\begin{tikzcd}
C \arrow[r, "x_1\lhd x_2"] & D \\
A \arrow[u, "x_1\rhd x_2"] \arrow[r, "x_1"'] \arrow[ur, phantom, "\circlearrowleft"] & B \arrow[u, "x_2"']
\end{tikzcd}$$

and a *corner trunk* is a trunk where the set of cubes satisfies the *bidistributivity* condition.

$$\begin{tikzcd}
 & G \arrow[rr, "{\begin{array}{c}(x_1 \lhd (x_2 \rhd x_3)) \lhd (x_2 \lhd x_3)\\ =\\ (x_1 \lhd x_2) \rhd x_3\end{array}}"] & & H \\
E \arrow[ur, "{\begin{array}{c}(x_1 \rhd x_2) \lhd ((x_1 \lhd x_2) \rhd x_3)\\ =\\ (x_1 \rhd x_2) \rhd (x_2 \lhd x_3)\end{array}}"] \arrow[rr, "x_1\lhd(x_2\rhd x_3)"'] & & F \arrow[ur, "x_2\lhd x_3"] & \\
 & C \arrow[uu, dotted, "(x_1\lhd x_2)\rhd x_3"] \arrow[rr, dotted, "x_1\lhd x_2"] & & D \arrow[uu, "x_3"'] \\
A \arrow[uu, "{\begin{array}{c}(x_1 \rhd x_2) \rhd (x_1 \rhd x_3)\\ =\\ x_1(\rhd x_2 \rhd x_3)\end{array}}"] \arrow[ur, dotted, "x_1\rhd x_2"'] \arrow[rr, "x_1"] & & B \arrow[uu, "x_2\rhd x_3"] \arrow[ur, "x_2"'] &
\end{tikzcd}$$

As a group can be seen as a category with one object and all its arrows invertible, to any rack X we can associate a corner trunk $X_{\mathbf{Tr}}$ $\big($Example (3.4)$\big)$. Moreover, as there is a (pre)simplicial set from a category (the nerve), we there is a *precubical set* $N(\mathcal{T})$ from a trunk $\mathcal{T}$, that is, a contravariant functor from the category $\square$ to the category **Set**, by $N(\mathcal{T})(-) := \mathrm{Hom}_{\mathbf{Trunk}}(-,\mathcal{T})$ $\big($cf. (6)$\big)$. In the case where the trunk is the trunk canonically associated to a rack, the description of the precubical nerve is easy to compute $\big($Theorem (3.9)$\big)$, and these result allows us to give a cubical description of the rack cohomology $\big($Corollary (3.11)$\big)$. Then we use this description to define our dendriform structure $\big($formulas (10) and (11)$\big)$. Our interest for this cubical definition is that the proof that the differential graded dendriform structure on $HR^{\bullet}(-)$ is well defined becomes a succession of combinatorial problems.

Section 4 : A graded dendriform algebra structure on $HR^\bullet(X,A)$. This section and the following are the heart of this paper. We use the cubical description of the rack cohomology made using trunks in the Section 3 to define a graded dendriform algebra structure on the cochain complex $\{CR^n(X,A), d_R^n\}_{n\in\mathbb{N}}$ computing the rack cohomology of a rack X with coefficients in a associative algebra (considered as a trivial X-module) (Theorem (4.2)). Then we show that this structure is compatible with the differential (Theorem (4.3)).

Dendriform and Zinbiel algebras are closely related to shuffles as we can see on examples (1.1) and (1.3). Then, in order to define such an algebraic structure on our cochain complex, we have to link shuffles and some sets of trunk maps. We make this possible by showing the existence of a map (9)

$$\rho : \mathrm{Sh}_{p_1,p_2} \to \mathrm{Hom}_{\mathbf{Set}}\big(N(X_{\mathbf{Tr}})(\square_{p_1+p_2}), N(X_{\mathbf{Tr}})(\square_{p_1}) \times N(X_{\mathbf{Tr}})(\square_{p_2})\big).$$

satisfying good properties with respect to the composition $\circ$ and the concatenation product $\star$ (2) defined on $\mathbb{S}_n$ (Lemma (4.1)).

Section 5 : A graded associative algebra morphism from $H^\bullet(G,A)$ to $HR^\bullet(\mathrm{Conf}(G),A)$. In this section we recall the definitions of the group cohomology $H^\bullet(-)$ and of the cup product defined on it. Seeing a group G as a category $G_{\mathbf{Cat}}$ and using the simplicial nerve $\mathrm{B}(G_{\mathbf{Cat}})$, we recall a simplicial definition of group cohomology (with trivial coefficient) and of its cup product. Now, on one hand we have a cohomology theory based on simplicial relations and on the other hand a cohomology theory based on cubical relations. Therefore, in order to define the expected morphism from $H^\bullet(G,A)$ to $HR^\bullet(\mathrm{Conf}(G),A)$, we have to find the good relation between simplices and cubes. The construction of this morphism $\big($Theorem (5.6)$\big)$ is based on the decomposition of the n-cubes into $n!$ n-simplex, that is, the existence of functors $\sigma : \Delta_n \to \square_n$ for all $\sigma \in \mathbb{S}_n$. We can represent geometrically such a functor in the following maneer $(n=3, \sigma=(13))$



We finish this section by proving that this morphism is compatible with the graded associative structures on $H^\bullet(-)$ and $HR^\bullet(-)$ $\big($Theorem (5.8)$\big)$.

1 Dendriform and Zinbiel algebras

Shuffles. Let $n \in \mathbb{N}$, we denote by $\mathbb{S}_n$ the group of permutations of $\{1,\dots,n\}$. Let $p_1, p_2 \in \mathbb{N}$, a (p_1,p_2)-*shuffle* is an element σ of $\mathbb{S}_{p_1+p_2}$ satisfying

$$\sigma(1) < \cdots < \sigma(p_1) \text{ and } \sigma(p_1+1) < \cdots < \sigma(p_1+p_2).$$

Remark that σ satisfies $\sigma(p_1) = p_1 + p_2$ or $\sigma(p_1 + p_2) = p_1 + p_2$.

Denote by Sh_{p_1,p_2} the set of (p_1, p_2)-shuffles, $\mathrm{Sh}^{p_1}_{p_1,p_2}$ the subset of elements $\sigma \in \mathrm{Sh}_{p_1,p_2}$ satisfying $\sigma(p_1) = p_1 + p_2$ and $\mathrm{Sh}^{p_1+p_2}_{p_1,p_2}$ the subset of elements $\sigma \in \mathrm{Sh}_{p_1,p_2}$ satisfying $\sigma(p_1 + p_2) = p_1 + p_2$.

Identifying a permutation $\sigma \in \mathbb{S}_n$ with the n-tuples $(\sigma(1), \ldots, \sigma(n))$, the set Sh_{p_1,p_2} corresponds to the set of n-tuples $(a_1, \ldots, a_{p_1+p_2})$ satisfying $a_1 < \cdots < a_{p_1}$ and $a_{p_1+1} < \cdots < a_{p_1+p_2}$. The cardinal of Sh_{p_1,p_2} (resp. $\mathrm{Sh}^{p_1}_{p_1,p_2}$, $\mathrm{Sh}^{p_1+p_2}_{p_1,p_2}$) is the binomial coefficient $\begin{pmatrix} p_1+p_2 \\ p_1 \end{pmatrix}$ (resp. $\begin{pmatrix} p_1+p_2 \\ p_1-1 \end{pmatrix}$, $\begin{pmatrix} p_1+p_2-1 \\ p_1 \end{pmatrix}$).

In the same way, given $p_1, p_2, p_3 \in \mathbb{N}$, the set of (p_1, p_2, p_3)-*shuffles* is defined as the subset of elements $\sigma \in \mathbb{S}_{p_1+p_2+p_3}$ satisfying

$$\sigma(1) < \cdots < \sigma(p_1)\,,\ \sigma(p_1+1) < \cdots < \sigma(p_1+p_2) \ \text{ and } \ \sigma(p_1+p_2+1) < \cdots < \sigma(p_1+p_2+p_3).$$

Let $p_1, p_2, p_3 \in \mathbb{N}$, there are bijections

$$\mathrm{Sh}_{p_1,p_2+p_3} \times \mathrm{Sh}_{p_2,p_3} \xrightarrow{\alpha} \mathrm{Sh}_{p_1,p_2,p_3} \ \text{ and } \mathrm{Sh}_{p_1+p_2,p_3} \times \mathrm{Sh}_{p_1,p_2} \xrightarrow{\beta} \mathrm{Sh}_{p_1,p_2,p_3} \tag{1}$$

given by $\alpha(\sigma, \gamma) = \sigma \circ (1_{p_1} \star \gamma)$ and $\beta(\sigma, \gamma) = \sigma \circ (\gamma \star 1_{p_3})$ where $\star : \mathbb{S}_p \times \mathbb{S}_q \to \mathbb{S}_{p+q}$ is the map defined by

$$(\sigma \star \gamma)(k) := \begin{cases} \sigma(k) & \text{if } 1 \le k \le p, \\ p + \gamma(k-p) & \text{if } p+1 \le k \le p+q. \end{cases} \tag{2}$$

In subset notation, formulas for α and β are

$$\alpha\big((a_1, \ldots, a_{p_1+p_2+p_3}), (b_1, \ldots, b_{p_2+p_3})\big) = (a_1, \ldots, a_{p_1}, a_{p_1+b_1}, \ldots, a_{p_1+b_{p_2+p_3}})$$

and

$$\beta\big((a_1, \ldots, a_{p_1+p_2+p_3}), (b_1, \ldots, b_{p_1+p_2})\big) = (a_{b_1}, \ldots, a_{b_{p_1+p_2}}, a_{p_1+p_2+1}, \ldots, a_{p_1+p_2+p_3}).$$

As we will see in the examples (1.1) and (1.3), the notion of a shuffle is closely related to the notion of a dendriform algebra.

Dendriform algebras. A (graded) dendriform algebra is a (graded) vector space D provided with two (graded) linear maps,

$$\succ, \prec : D \otimes D \to D,$$

which satisfy for all $x, y, z \in D$ the following relations

$$\begin{aligned} &x \succ (y \succ z) = (x \succ y) \succ z + (x \prec y) \succ z \\ &(x \succ y) \prec z = x \succ (y \prec z) \\ &(x \prec y) \prec z = x \prec (y \prec z) + x \prec (y \succ z) \end{aligned}$$

Example 1.1. $T(V)$, free dendriform algebra

- Let V be a vector space, we define a dendriform algebra structure on $T(V)$ by putting

$$v_1 \ldots v_p \succ v_{p+1} \ldots v_{p+q} := \sum_{\sigma \in Sh^{p+q}_{p,q}} v_{\sigma^{-1}(1)} \ldots v_{\sigma^{-1}(p+q-1)} v_{p+q},$$

and

$$v_1 \dots v_p \prec v_{p+1} \dots v_{p+q} := \sum_{\sigma \in Sh^p_{p,q}} v_{\sigma^{-1}(1)} \dots v_{\sigma^{-1}(p-1)} v_{p+q} v_{\sigma^{-1}(p+1)} \dots v_{\sigma^{-1}(p+q)}.$$

- Let V be a vector space, J.-L. Loday has defined a structure of dendriform algebra on $\bigoplus_{n\geq 0} \mathbb{K}[Y_n] \otimes V^{\otimes n}$ where Y_n is the set of planar binary trees with $n+1$ leaves (cf. [Lod01]). This algebra is the free dendriform algebra associated to V.
- This example is an example of graded dendriform algebra using the well known *Pascal formula.*

  Let A be a associative algebra and consider the graded $\mathbb{K}$-module $\bigoplus_{n\geq 0} A_n$ where $A_n = A$ for all $n \in \mathbb{N}$. We define a structure of graded dendriform algebra on it by putting for $a \in A_p$ and $b \in A_q$

  $$a \succ b := \binom{p+q-1}{p} ab,$$

  and

  $$a \prec b := \binom{p+q-1}{p-1} ab.$$

Relation with associative algebras. A (graded) dendriform algebra is a particular example of (graded) associative algebra. Indeed, let $(D, \succ, \prec)$ be a (graded) dendriform algebra, we define a new product $*$ on D by putting

$$x * y := x \succ y + x \prec y.$$

Proposition 1.2. *The product $*$ is associative.*

Hence there exists a functor from the category of dendriform algebras to the category of associative algebras

$$\mathbf{Dend} \to \mathbf{As}$$

The product $*$ is non necessarily commutative but it becomes if $x \succ y = y \prec x$. This condition leads us to the notion of *commutative dendriform algebra*, which is also called *Zinbiel algebra.*

Zinbiel algebras. A Zinbiel algebra is a vector space D provided with a linear map

$$\succ : D \otimes D \to D,$$

which satisfies for all $x, y, z \in D$ the following relation

$$x \succ (y \succ z) = (x \succ y) \succ z + (y \succ x) \succ z.$$

Remark that the variables do not stay in the same order, thus in the graded case we have to be careful. A *graded Zinbiel algebra* is a graded vector space D provided with a graded linear map $\succ : D \otimes D \to D$ which staisfies for all $x, y, z \in D$ the relation

$$x \succ (y \succ z) = (x \succ y) \succ z + (-1)^{pq} (y \succ x) \succ z,$$

where q is the degree of y.

Example 1.3. $T(V)$, $HL^{\bullet}(\mathfrak{g}, A)$

- Let V be a vector space, there is a structure of Zinbiel algebra defined on $T(V)$ by putting

  $$v_1 \dots v_p \succ v_{p+1} \dots v_{p+q} := \sum_{\sigma \in Sh_{p,q}^{p+q}} v_{\sigma^{-1}(1)} \dots v_{\sigma^{-1}(p+q-1)} v_{p+q}.$$

  There is a structure of graded Zinbiel algebra definedon $T(V)$ by

  $$v_1 \dots v_p \succ v_{p+1} \dots v_{p+q} := \sum_{\sigma \in Sh_{p,q}^{p+q}} \epsilon(\sigma) v_{\sigma^{-1}(1)} \dots v_{\sigma^{-1}(p+q-1)} v_{p+q}. \tag{3}$$

- Let $\mathfrak{g}$ be a Leibniz algebra and A a commutative algebra. In [Lod95] J.-L. Loday has defined a graded Zinbiel product on $HL^*(\mathfrak{g}, A)$, the Leibniz cohomology of $\mathfrak{g}$ with values in A see as a trivial $\mathfrak{g}$-representation.

Relation with dendriform algebras and commutative algebras. As it said above, a Zinbiel algebra is a kind of commutative dendriform algebra. Indeed, let $(D, \succ)$ be a Zinbiel algebra, and define a second product $\prec$ by the formula $x \prec y := y \succ x$ for all $x, y \in D$.

Proposition 1.4. *$(D, \succ, \prec)$ is a dendriform algebra, and $(D, *)$ is commutative. Conversely, a dendriform algebra $(D, \succ, \prec)$ where $x \succ y = y \prec x$ for all $x, y \in D$, is a Zinbiel algebra.*

Hence, there exist functors from the category of Zinbiel algebras to the category of commutative algebras, and to the category of dendriform algebras. All these functors fit into the following diagram

$$\begin{tikzcd}
& \mathbf{Dend} \arrow[dr] & \\
\mathbf{Zinb} \arrow[ur] \arrow[dr] & & \mathbf{As} \\
& \mathbf{Com} \arrow[ur] &
\end{tikzcd}$$

Remark 1.5. This diagram fits into a (beautiful) bigger diagram which is (cf. [Lod01])

$$\begin{tikzcd}
& \mathbf{Dend} \arrow[dr] & & \mathbf{Dialg} \arrow[dr] & \\
\mathbf{Zinb} \arrow[ur] \arrow[dr] & & \mathbf{As} \arrow[ur] \arrow[dr] & & \mathbf{Leib} \\
& \mathbf{Com} \arrow[ur] & & \mathbf{Lie} \arrow[ur] &
\end{tikzcd}$$

where **Dialg** is the category of dialgebras, **Leib** is the category of Leibniz algebras, and **Lie** is the category of Lie algebras. We can remark that each operad on the left (**Zinb, Dend, Com, As**) is Koszul dual to the operad on the right (**Leib, Dialg, Lie, As**) which is symetric to it with respect to the vertical line passing through **As**.

2 Racks

Shelves. A *shelf* is a set X provided with a binary product $\rhd : X \times X \to X$ satisfying $x \rhd (y \rhd z) = (x \rhd y) \rhd (x \rhd z)$ for all $x, y, z \in X$. A *pointed shelf* is the data of a shelf $(X, \rhd)$ provided with an element $1 \in X$, called the unit, satisfying $1 \rhd x = x$ and $x \rhd 1 = 1$ for all $x \in X$.

Notation : Let $(X, \rhd)$ be a shelf. For all $x \in X$ denote by c_x the map from X to X defined by $c_x(y) = y$. Because in a shelf the product $\rhd$ is non associative we have to be careful with the parenthesis. In the sequel the expression $(c_{x_1} \circ \cdots \circ c_{x_{n_1}})(x_n)$ (bracketing by the right) is denoted by $x_1 \rhd \ldots \rhd x_n$.

Let X and X' be two shelves. A *morphism of shelves* from X to X' is a map $f : X \to X'$ which preserves the products. If the shelves are pointed then a *morphism of pointed shelves* is a morphism of shelves which preserves the units.

The category of shelves (resp. pointed shelves) is denoted by **Shelf** (resp. **pShelf**).

Racks. A *(pointed) rack* is a (pointed) shelf $(X, \rhd)$ where for all $x \in X$ the map $c_x : X \to X$ is a bijection. Let X and X' two (pointed) racks. A morphism of (pointed) racks from X to X' is a morphism of (pointed) shelves $f : X \to X'$.

The category of racks (resp. pointed racks) is denoted by **Rack** (resp. **pRack**).

Example 2.1. Groups and augmented racks

- Let G be a group. Define a rack on the set G by taking $x \rhd y = xyx^{-1}$. This rack is denoted $Conj(G)$. It is pointed by e the unit of G.
- Let G be a group, X be a G-set and $f : X \to G$ a G-map (where G acts on itself by connjugation). Define a rack structure on X by taking $x \rhd y = f(x) \cdot y$. If there exists a fixed point $1 \in X$ such that $f(1) = e$, then this rack is pointed by 1.

In Example (2.1) a functor Conj : **Grp** → **Rack** from the category of groups to the category of racks is defined. There is a functor As : **Rack** → **Grp** defined by

$$\begin{aligned} \mathrm{As}(X) &:= \mathcal{F}(X)/ < x \rhd y = xyx^{-1} >, \\ \mathrm{As}(f) &:= \overline{\mathcal{F}(f)}, \end{aligned}$$

where $\mathcal{F}(X)$ is the free group generated by the set X.

Proposition 2.2. *The functor* As *is left adjoint to the functor* Conj.

Rack modules. Let X be a rack. A *left X-module* is an abelian group A provided with a map

$$\cdot : X \times A \to A; (x, a) \mapsto x \cdot a$$

satisfying $x \cdot (y \cdot a) = (x \rhd y) \cdot (x \cdot a)$ for all $a \in A$ and $x, y \in X$. If the rack is pointed then we ask for the additional axiom : $1 \cdot a = a, \forall a \in A$. Let A and B be two left X-module. A map $f : A \to B$ is a morphism of left X-module if $f(x \cdot a) = x \cdot f(a)$ for all $(a, x) \in A \times X$. We denote by $X - \mathbf{Mod}$ the category of left X-module.

Example 2.3. Let G be a group and A be a left G-module, then A is a left Conj(G)-module.

Remark 2.4. Let X be a rack and A be a left X-module. There is an equivalence of categories between $X - \mathbf{Mod}$ and $\mathrm{As}(X) - \mathbf{Mod}$.

Rack cohomology. Let X be a rack and A be a left X-module. We define a cochain complex $\{CR^n(X,A), d_R^n\}_{n\in\mathbb{N}}$ by

$$CR^n(X,A) := \mathrm{Hom}_{\mathbf{Set}}(X^n, A),$$
$$d_R^{n+1} f := \sum_{i=1}^{n+1} (-1)^i \big(d_{i,0}^{n+1} f - d_{i,1}^{n+1} f\big),$$

where

$$d_{i,\epsilon}^{n+1} f(x_1,\dots,x_{n+1}) := \begin{cases} f(x_1,\dots,x_{i-1},x_i \triangleright x_{i+1},\dots,x_i \triangleright x_{n+1}) & \text{if } \epsilon = 0, \\ (x_1 \triangleright \dots \triangleright x_i)\cdot f(x_1,\dots,x_{i-1},x_{i+1},\dots,x_{n+1}) & \text{if } \epsilon = 1. \end{cases}$$

The fact that d_R is a differential comes from the *cubical identities* satisfied by the family of maps $\{d_{i,\epsilon}^n\}$, that is, for all $1 \le i < j \le n+1$ and $\epsilon,\omega \in \{0,1\}$, we have the identities :

$$d_{i,\epsilon}^{n+1} \circ d_{j-1,\omega}^{n} = d_{j,\omega}^{n+1} \circ d_{i,\epsilon}^{n}.$$

The cohomology associated to this cochain complex is called the *rack cohomology of X with coefficients in A* and denoted $HR^\bullet(X,A)$

Remark 2.5. In this paper we will consider the rack cohomology with trivial coefficients, that is when A is a trivial left X-module. In this case we have $d_{n+1,0}^{n+1} f = d_{n+1,1}^{n+1} f$ for all $f \in CR^n(X,A)$.

Remark 2.6. If X is a pointed rack, then a new cochain complex $CR_p^\bullet(X,A) = \{CR_p^n(X,A), d_R^n\}_{n\ge 1}$ is defined by

$$CR_p^n(X,A) := \{f \in \mathrm{Hom}_{\mathbf{Set}}(X^n,A) \,|\, f(x_1,\dots,1,\dots,x_n) = 0\}.$$

The cohomology associated to this cochain complex is called the *pointed rack cohomology of X with coefficients in A* and denoted $HR_p^\bullet(X,A)$.

Remark 2.7. In the definition of $\{CR^n(X,A), d_R^n\}_{n\in\mathbb{N}}$ (resp. $\{CR_p^n(X,A), d_R^n\}_{n\in\mathbb{N}}$ we don't use the second axiom of a rack, that is the bijectivity of c_x for all $x \in X$. Hence this cochain complex is well defined for a shelf (resp. pointed shelf). It correponds to the cochain complex associated to the *multi-shelf* $(X,\triangleright,\triangleleft)$, where $x \triangleleft y = x$, defined in [PS].

3 The category of trunks

Trunks are the fundamental tools in order to solve our problems combinatorially. This section is based on [FRS95]. Note that our definitions are slightly different for we are working in the left rack context instead of the right rack context. The two theories are obviously equivalent.

3.1 Definitions and examples

Categories. Following [Mac98] let us recall the definition of a (small) category as a graph provided with extra properties. Let $\mathcal{G} = (V,E,s,t)$ be a directed graph with V the set of vertices, E the set of edges, $s : E \to V$ the source map and $t : E \to V$ the target map. We define a subset of $E \times E \times E$ by

$$E \times_V E \times_V E := \{(a,b,c) \in E \times E \times E \,|\, t(a) = s(b),\ t(b) = s(c)\},$$

and subsets of $E \times E$ by

$$\begin{aligned} E \times_V E &:= \{(a,b) \in E \times E \,|\, t(a) = s(b)\}, \\ V \times_V E &:= \{(s(a),a) \in E \times E \,|\, a \in E\}, \\ E \times_V V &:= \{(a,t(a)) \in E \times E \,|\, a \in E\}. \end{aligned}$$

A *category* is a directed graph $\mathcal{C} = (V, E, s, t)$ provided with a *composition map* $c : E \times_V E \to E$ and an *identity map* $i : V \to E$ satisfying the following commutative diagrams.

$$\begin{tikzcd} E \times_V E \times_V E \arrow[r, "c\times 1"] \arrow[d, "1\times c"'] & E \times_V E \arrow[d, "c"] \\ E \times_V E \arrow[r, "c"] & E \end{tikzcd} \qquad \begin{tikzcd} V \times_V E \arrow[r, "i\times 1"] \arrow[dr, "\pi_2"] & E \times_V E \arrow[d, "c"] & E \times_V V \arrow[l, "1\times i"'] \arrow[dl, "\pi_1"'] \\ & E & \end{tikzcd}$$

The left diagram states the *associativity* of the composition c and the right diagram states that each $i(x)$ is a *left and right unit* for the composition.

Example 3.1. Simplexes, cubes and groups.

- Let $n \in \mathbb{N}$, let us consider the directed graph $\Delta_n = \{V(\Delta_n), E(\Delta_n), s, t\}$ where $V(\Delta_n) := \{0, \ldots, n\}$, $E(\Delta_n) := \{(i,j) \in V(\Delta_n) \times V(\Delta_n) \,|\, i \leq j\}$, $s(i,j) = i$ and $t(i,j) = j$. Define a composition map c by $c\big((i,j),(j,k)\big) = (i,k)$ and an identity map i by $i(j) = (j,j)$.
- Let $n \in \mathbb{N}$, let us consider the directed graph $\square_n = \{V(\square_n), E(\square_n), s, t\}$ where $V(\square_n) := \{A \subseteq \{1, \ldots, n\}\}$, $E(\square_n) := \{(A,B) \in V(\square_n) \times V(\square_n) \,|\, A \subseteq B\}, s(A,B) = A$ and $t(A,B) = B$. Define a composition map c by $c\big((A,B),(B,C)\big) = (A,C)$ and an identity map i by $i(A) = (A,A)$.
- Let G be a group. Consider the directed graph $G = (\{\star\}, G, s, t)$ with $s = t : G \to \{\star\}$, the canonical map from G to the terminal object in **Set**. Define a composition map $c : G \times G \to G$ by $c(a,b) = \mu(a,b)$ (the product of a and b in G) and an identity map $i : \{\star\} \to G$ by $i(\star) = e$ (the unit in G).

Functors. Let $\mathcal{C}$ and $\mathcal{C}'$ be two categories. A *functor* from $\mathcal{C}$ to $\mathcal{C}'$ is a graph map $F : \mathcal{C} \to \mathcal{C}'$ which preserves the composition and the units, i.e F satisfies the following commutative diagrams

$$\begin{tikzcd} E \times_V E \arrow[r, "c"] \arrow[d, "F\times F"'] & E \arrow[d, "F"] \\ E' \times_V E' \arrow[r, "c'"] & E' \end{tikzcd} \qquad \begin{tikzcd} V \arrow[r, "i"] \arrow[d, "F"'] & E \arrow[d, "F"] \\ V' \arrow[r, "i'"] & E' \end{tikzcd}$$

Example 3.2. Faces of simplexes, faces of cubes and permutations.

- Let $n \in \mathbb{N}$. For all $0 \leq i \leq n$ define a functor ∂_i^n from the graph Δ_{n-1} to Δ_n by

  $$\partial_i^n(k) := \begin{cases} k & \text{if } 0 \leq k \leq i-1, \\ k+1 & \text{if } i \leq k \leq n-1. \end{cases}$$

  Given two vertices in Δ_{n-1} there is exactly one edge between them, hence there is only one way to define ∂_i^n on the set of edges in order to have a graph map.

- Let $n \in \mathbb{N}$. For all $(i,\epsilon) \in \{1,\dots,n\} \times \{0,1\}$ define a functor $\partial^n_{i,\epsilon}$ from the graph $\Box_{n-1}$ to $\Box_n$ by
$$\partial^n_{i,\epsilon}(A) = \left\{ \begin{array}{ll} A_{<i} \amalg t_{+1}(A_{\geq i}) & \text{if } \epsilon = 0, \\ A_{<i} \amalg t_{+1}(A_{\geq i}) \amalg \{i\} & \text{if } \epsilon = 1. \end{array} \right.$$
where $A_{<i} = \{a \in A \,|\, a < i\}$, $A_{\geq i} = \{a \in A \,|\, a \geq i\}$ and $t_{+1}(B) = \{b+1 \,|\, b \in B\}$. Given two vertices in $\Box_{n-1}$ there is at most one edges between them, hence there is only one way to define $\partial^n_{i,\epsilon}$ on the set of edges in order to have a graph map.

- Let $n \in \mathbb{N}$. For all $\sigma \in \mathbb{S}_n$ define a functor σ from the graph Δ_n to the graph $\Box_n$ by
$$\sigma(0) := \emptyset \text{ and } \sigma(k) := \{\sigma(1), \dots, \sigma(k)\}\, \forall k \geq 1.$$

As explain in the motivations, we want to change the associativity relation by the *distributivity* relation.

Trunks. Let $\mathcal{G} = (V,E,s,t)$ be a directed graph. We denote by $\mathrm{S}(\mathcal{G})$ the subset of $E \times_V E \times E \times_V E$ of elements (a,b,c,d) satisfying $s(a) = s(c)$, $t(a) = s(b)$, $t(c) = s(d)$ and $t(b) = t(d)$. A *trunk* is a directed graph $\mathcal{T} = (V,E,s,t)$ provided with a subset Γ of $\mathrm{S}(\mathcal{T})$. An element in Γ is called a *preferred square.*

A *pointed trunk* is a trunk $(\mathcal{T} = (V,E,s,t),\Gamma)$ together with an identity map $i : V \to E$ satisfying $\big(a, i(t(a)), i(s(a)), a\big) \in \Gamma$ and $\big(i(s(a)), a, a, i(t(a))\big) \in \Gamma$ for all $a \in E$.

Example 3.3. Categories and cubes.

- Let $\mathcal{C} = (V,E,s,t)$ be a category. We define a trunk $\mathrm{Tr}(\mathcal{C})$ by taking the pair $(\mathcal{C},\Gamma)$ with Γ the set of commutative diagrams in $\mathcal{C}$
$$\Gamma = \{(a,b,c,d) \in \mathrm{S}(E) \,|\, c(a,b) = c(c,d)\}.$$
The identity map in the category $\mathcal{C}$ provides $\mathrm{Tr}(\mathcal{C})$ with a pointed trunk structure.

- Let $n \in \mathbb{N}$. Consider the graph $\Box_n$ with set of vertices $V(\Box_n) = \{A \subseteq \{1,\dots,n\}\}$ and set of edges $E(\Box_n) = \{(A, A \amalg \{k\}) \,|\, k \notin A\}$. Take as set of preferred square $\Gamma_{\Box_n}$ the set equals to
$$\{\big((A, A \amalg \{k\}), (A \amalg \{k\}, A \amalg \{k,l\}), (A, A \amalg \{l\}), (A \amalg \{l\}, A \amalg \{k,l\})\big) \,|\, k < l\}.$$

Corner trunks. A *corner trunk* is a trunk $(\mathcal{T},\Gamma)$ where Γ is the graph of a map $c : E \times_V E \to E \times_V E$ satisfying the following commutative diagram

$$\begin{tikzcd}
E \times_V E \times_V E \arrow[d, "c\times 1"'] & E \times_V E \times_V E \arrow[l, "1\times c"'] \arrow[r, "c\times 1"] & E \times_V E \times_V E \arrow[d, "1\times c"] \\
E \times_V E \times_V E \arrow[r, "1\times c"] & E \times_V E \times_V E & E \times_V E \times_V E \arrow[l, "c\times 1"']
\end{tikzcd}$$

The map c is called the *composition map.* The relation satisfied by c can be decomposed in three relations, called *bidistributivity relations.* To describe these relations, let us denote by $\rhd$ (resp. $\lhd$) the composition $\mathrm{pr}_1 \circ c$ (resp. $\mathrm{pr}_2 \circ c$). This defines maps $\rhd$ and $\lhd$ from $E \times_V E$ to E, and the relations are :

- $a \rhd (b \rhd c) = (a \rhd b) \rhd ((a \lhd b) \rhd c)$,

- $(a \rhd b) \lhd ((a \lhd b) \rhd c) = (a \lhd (b \rhd c)) \rhd (b \lhd c)$,
- $(a \lhd b) \lhd c = (a \lhd (b \rhd c)) \lhd (b \lhd c)$,

for all $(a, b, c) \in E \times_V E \times_V E$.

In the case where the trunk is pointed then we have necessarily the following relations, called *unit relations* :

- $a \rhd i(t(a)) = i(s(a))$ and $i(s(a) \rhd a = a$,
- $a \lhd i(t(a)) = a$ and $i(s(a)) \lhd a = i(t(a))$.

Example 3.4. Racks and $\mathbb{N}$.

- Let $(X, \rhd)$ be a rack or shelf. Consider the directed graph $\mathcal{T}_X = (\{\star\}, X, s, t)$ with $s = t : X \to \{\star\}$, the unique map from X to the terminal object in **Set**. Define a composition map $c : X^2 \to X^2$ by $c(a, b) = (a \rhd b, a)$. The set of preferred square Γ_X is equal to $\mathrm{S}(X) = X^4$. In this case, because the map $\lhd$ is the first projection, the second and third bidistributivity relations are trivials and the first is equivalent to the distributivity of $\rhd$. Remark that $\mathcal{T}_X$ is a pointed corner trunk if and only if X is a pointed rack/shelf.
- Consider the graph $\mathbb{N} = (\{\star\}, \mathbb{N}, s, t)$ with $s = t : \mathbb{N} \to \{\star\}$, the unique map from $\mathbb{N}$ to the terminal object in **Set**. Define a composition map $c : \mathbb{N}^2 \to \mathbb{N}^2$ by $c(i, j) = (\max(i, j), \min(i, j))$. The set of preferred square $\Gamma_{\mathbb{N}}$ equals to $\mathrm{S}(\mathbb{N}) = \mathbb{N}^4$.

Trunk maps. Let $(\mathcal{T}, \Gamma)$ and $(\mathcal{T}', \Gamma')$ be two trunks. A *trunk map* from $\mathcal{T}$ to $\mathcal{T}'$ is a graph map $F : \mathcal{T} \to \mathcal{T}'$ mapping Γ to Γ'. If the trunks are pointed then F is a *pointed trunk map* if F preserves the units. In the case where the trunks $\mathcal{T}$ and $\mathcal{T}'$ are corner trunks then $F : \mathcal{T} \to \mathcal{T}'$ is a trunk map if and only if F preserves the composition map c, or, if and only if F preserves the products $\rhd$ and $\lhd$.

Example 3.5. Functors, faces of a cube and shuffles.

- Let F be a functor from a category $\mathcal{C}$ to a category $\mathcal{C}'$. Because F maps a commutative diagram in $\mathcal{C}$ to a commutative diagram in $\mathcal{C}'$, it induces a pointed trunk map $\mathrm{Tr}(F)$ from the pointed trunk $\mathrm{Tr}(\mathcal{C})$ to $\mathrm{Tr}(\mathcal{C})$ (cf. Example (3.3)).
- Let $n \in \mathbb{N}$ and consider the graph map $\partial^n_{i,\epsilon} : \square_{n-1} \to \square_n$ defined in Example (3.2).

  Given $1 \leq k < l \leq n-1$, we have $\partial^n_{i,0}(\{k\}) < \partial^n_{i,0}(\{l\})$. Moreover for all A, B such that $A \cap B = \emptyset$ we have $\partial^n_{i,\epsilon}(A \amalg B) = \partial^n_{i,\epsilon}(A) \amalg \partial^n_{i,0}(B)$. Thus $\partial^n_{i,\epsilon}$ maps a preferred square in $\square_{n-1}$ to a preferred square in $\square_n$.
- Let $p_1, p_2 \in \mathbb{N}$ and $\sigma \in \mathrm{Sh}_{p_1,p_2}$. Let us define from σ a graph map $\sigma : \square_{p_1+p_2} \to \square_{p_1+p_2}$ by

  $$\sigma(\{a_1, \dots, a_k\}) := \{\sigma(a_1), \dots, \sigma(a_k)\}.$$

  Define two graph maps $\sigma \circ i_{p_1} : \square_{p_1} \to \square_{p_1+p_2}$ and $\sigma \circ i_{p_2} : \square_{p_2} \to \square_{p_1+p_2}$ where

  $$\begin{aligned} i_{p_1} &:= \partial^{p_1+p_2}_{p_1+p_2,0} \circ \dots \circ \partial^{p_1+1}_{p_1+1,0}, \\ i_{p_2} &:= \partial^{p_1+p_2}_{1,1} \circ \dots \circ \partial^{p_2+1}_{1,1}. \end{aligned}$$

  We have

  $$\begin{aligned} (\sigma \circ i_{p_1})(A) &= \{\sigma(a_1), \dots, \sigma(a_k)\}, \\ (\sigma \circ i_{p_2})(A) &= \{\sigma(1), \dots, \sigma(p), \sigma(p + a_1), \dots, \sigma(p + a_k)\}. \end{aligned}$$

Let $1 \leq k < l \leq p_1$, $\sigma \in \mathrm{Sh}_{p_1,p_2}$ implies $\sigma(k) < \sigma(l)$. Moreover for all A, B such that $A \cap B = \emptyset$ we have $(\sigma \circ i_{p_1})(A \amalg B) = (\sigma \circ i_{p_1})(A) \amalg (\sigma \circ i_{p_1})(B)$, thus $(\sigma \circ i_{p_1})(A \amalg \{k\}) = (\sigma \circ i_{p_1})(A) \amalg \{\sigma(k)\}$ and $(\sigma \circ i_{p_1})(A \amalg \{l\}) = (\sigma \circ i_{p_1})(A) \amalg \{\sigma(l)\}$ with $\sigma(k) < \sigma(l)$. Thus $\sigma \circ i_{p_1}$ maps a preferred square in $\square_{p_1}$ to a preferred square in $\square_{p_1+p_2}$ and $\sigma \circ i_{p_1} : \square_{p_1} \to \square_{p_1+p_2}$ is a trunk map.

In the same way, let $1 \leq k < l \leq p_2$, $\sigma \in \mathrm{Sh}_{p_1,p_2}$ implies $\sigma(p_1 + k) < \sigma(p_1 + l)$. Moreover for all A, B such that $A \cap B = \emptyset$ we have $(\sigma \circ i_{p_2})(A \amalg B) = (\sigma \circ i_{p_2})(A) \amalg (\sigma \circ i_{p_2})(B)$, thus $(\sigma \circ i_{p_2})(A \amalg \{k\}) = (\sigma \circ i_{p_2})(A) \amalg \{\sigma(p_1+k)\}$ and $(\sigma \circ i_{p_2})(A \amalg \{l\}) = (\sigma \circ i_{p_2})(A) \amalg \{\sigma(p_1+l)\}$ with $\sigma(p_1+k) < \sigma(p_1+l)$. Thus $\sigma_\circ i_{p_2}$ maps a preferred square in $\square_{p_2}$ to a preferred square in $\square_{p_1+p_2}$ and $\sigma \circ i_{p_2} : \square_{p_2} \to \square_{p_1+p_2}$ is a trunk map.

Using these maps, we will define the graded dendriform algebra product on rack/shelf cohomology.

The category of (pointed) trunks is the category with the (pointed) trunks as vertices and the (pointed) trunk maps as edges. This category is denoted by **Trunk** (resp. **pTrunk**).

Relations between categories and trunks. In Example (3.3) we have seen that from a category $\mathcal{C}$ we can define a trunk $\mathrm{Tr}(\mathcal{C})$. Moreover we have seen in Example (3.5) that a functor F from a category $\mathcal{C}$ to a category $\mathcal{C}'$ determines a trunk map $\mathrm{Tr}(F)$ from $\mathrm{Tr}(\mathcal{C})$ to $\mathrm{Tr}(\mathcal{C}')$. Hence a functor $\mathrm{Tr} : \mathbf{Cat} \to \mathbf{Trunk}$, from the category of categories **Cat** to the category of trunks **Trunk** is defined by

$$\begin{aligned} \mathbf{Cat} &\xrightarrow{\mathrm{Tr}} \mathbf{Trunk} \\ \mathcal{C} &\longmapsto \mathrm{Tr}(\mathcal{C}) \\ F &\longmapsto \mathrm{Tr}(F) \end{aligned}$$

Let $(\mathcal{T}, \Gamma)$ be a trunk. Consider the free category $\mathcal{F}(\mathcal{T})$ generated by the graph $\mathcal{T}$. For all pair of objects (x, y) in $\mathcal{F}(\mathcal{T})$ we define a binary relation $R_{x,y}$ on $\mathrm{Hom}_{\mathcal{F}(\mathcal{T})}(x, y)$ by $(a_1, \ldots, a_n) R_{x,y} (b_1, \ldots, b_m)$ if and only the two following points are satisfied :

1. $n = m$,
2. For all $1 \leq k \leq n$ such that $a_k \neq b_k$ then $(a_k, a_{k+1}, b_k, b_{k+1}) \in \Gamma$ or $(a_{k-1}, a_k, b_{k-1}, b_k) \in \Gamma$.

Then we define a category $\mathrm{Cat}(\mathcal{T})$ by taking the quotient of $\mathcal{F}(\mathcal{T})$ by the binary relation R (cf. [Mac98] for the definition of a quotient category). Then the category $\mathrm{Cat}(\mathcal{T})$ has the same vertices as $\mathcal{T}$ and the set of edges is the set of strings of composable edges in $\mathcal{T}$ where two strings of edges are identified if they have the same length and are equal "up to preferred squares in Γ". Hence all commutative diagrams in $\mathrm{Cat}(\mathcal{T})$ comes from a preferred square in $(\mathcal{T}, \Gamma)$.

$$x_0 \overset{a_0}{\to} x_1 \overset{a_1}{\to} \cdots \overset{a_n}{\to} x_{n+1} \sim x_0 \overset{b_0}{\to} x_1 \overset{b_1}{\to} \cdots \overset{b_n}{\to} x_{n+1}$$

if and only if

$$\begin{tikzcd}
 & & x_2 \arrow[dr, "a_2"] & & & & x_{n-2} \arrow[dr, "a_{n-2}"] & & & \\
x_0 \arrow[r, "a_0"] & x_1 \arrow[ur, "a_1"] \arrow[dr, "b_1"'] & \in \Gamma & x_3 \arrow[r, "a_3"] & \ldots \arrow[r, "a_{n_4}"] & x_{n-3} \arrow[ur, "a_{n-3}"] \arrow[dr, "b_{n-3}"'] & \in \Gamma & x_{n-1} \arrow[r, "a_{n-1}"] & x_n \arrow[r, "a_n"] & x_{n+1} \\
 & & y_2 \arrow[ur, "b_2"'] & & & & y_{n-2} \arrow[ur, "b_{n-2}"'] & & &
\end{tikzcd}$$

Let F be a trunk map from $(\mathcal{T},\Gamma)$ to $(\mathcal{T}',\Gamma')$. Because F is a graph map, $\mathcal{F}(F)$ is a functor from $\mathcal{F}(\mathcal{T})$ to $\mathcal{F}(\mathcal{T}')$. Because F maps preferred square in Γ to preferred square in Γ', $\mathrm{Cat}(F)$ is a well defined functor from $\mathrm{Cat}(\mathcal{T})$ to $\mathrm{Cat}(\mathcal{T}')$.

Hence, there is a functor Cat : **Trunk** → **Cat**, from the category of trunks to the category of categories, defined by

$$\begin{aligned} \mathbf{Trunk} &\xrightarrow{\mathrm{Cat}} \mathbf{Cat} \\ \mathcal{T} &\longmapsto \mathrm{Cat}(\mathcal{T}) \\ F &\longmapsto \mathrm{Cat}(F) \end{aligned}$$

and there is an adjunction

$$\mathrm{Cat} \overset{\theta}{\vdash} \mathrm{Tr}. \tag{4}$$

3.2 Nerve of a trunk

The category $\square$. Objects of the *the cubical category* $\square$ are the graphs $\square_n$ defined in Example (3.1). Given $n \in \mathbb{N}$ vertices of $\square_n$ has been defined as the subsets of $\{1,\dots,n\}$. The bijection between the set of subsets of $\{1,\dots,n\}$ and the set $\{0,1\}^n$, given by $A \mapsto (\epsilon_1,\dots,\epsilon_n)$ with $\epsilon_k = 1$ if $k \in A$ and 0 if not, allows us to describe $\square_n$ in a second way. In the first case we said that $\square_n$ is described in subset notation, and in the second case in coordinate notation.

For all $m, n \in \mathbb{N}$, define $\mathrm{Hom}_\square(\square_m,\square_n)$ as the subset of $\mathrm{Hom}_{\mathbf{Graph}}(\square_m,\square_n)$ generated by the face maps $\partial^k_{i,\epsilon}$ defined in Example (3.5). In other words a morphism $\mathrm{Hom}_\square(\square_m,\square_n)$ is a graph map from $\square_m$ to $\square_n$ which is a composition of face maps. Using the description of $\square_n$ in term of coordinates, $\partial^n_{i,\epsilon}$ is the graph maps defined by

$$\partial^n_{i,\epsilon}(\epsilon_1,\dots,\epsilon_{n-1}) = (\epsilon_1,\dots,\epsilon_{i-1},\epsilon,\epsilon_i,\dots,\epsilon_{n-1}).$$

The next proposition states that face maps satisfy relations called *cubical relations*. This allows us to write a morphism in $\mathrm{Hom}_\square(\square_m,\square_n)$ in a canonical way.

Proposition 3.6 (Cubical relations)**.** *For all $1 \leq i < j \leq k+1$, and $\epsilon,\omega \in \{0,1\}$ we have the following relation :*

$$\partial^{k+1}_{i,\epsilon} \circ \partial^k_{j-1,\omega} = \partial^{k+1}_{j,\omega} \circ \partial^k_{i,\epsilon}. \tag{5}$$

Corollary 3.7. *Let $m < n \in \mathbb{N}$. Each element f in $\mathrm{Hom}_\square(\square_m,\square_n)$ can be written uniquely as*

$$f = \partial^n_{i_{n-m},\epsilon_{n-m}} \circ \cdots \circ \partial^{m+1}_{i_1,\epsilon_1},$$

with $i_1 < i_2 < \cdots < i_{n-m}$.

Cubical object in a category. Let $\mathcal{C}$ be a category. A *precubical object in $\mathcal{C}$* is a functor $N : \square^{op} \to \mathcal{C}$. Dually a *precocubical object in $\mathcal{C}$* is a functor $N : \square \to \mathcal{C}$.

Let $\mathbb{K}$ be a commutative ring and $\mathcal{C}$ be the category $\mathbf{Mod}_{\mathbb{K}}$ of modules over $\mathbb{K}$. From a precocubical $\mathbb{K}$-module M (that is a precocubical object in $\mathbf{Mod}_{\mathbb{K}}$), define a cochain complex $M^\bullet = \{M^n, d^n\}$ by

$$M^n := M(\square_n)\,,\; d^n := \sum_{i=1}^{n}(-1)^i(M(\partial^n_{i,0}) - M(\partial^n_{i,1})),$$

The graded map d^n is a differential thanks to the cubical relations (Proposition (3.6)).

Remark 3.8. Let A be a module over $\mathbb{K}$. From a precubical set $S : \square \to \mathbf{Set}$ we can always define a precocubical $\mathbb{K}$-module by postcomposed S by the functor $\mathrm{Hom}_{\mathbf{Set}}(-, A)$.

In the following paragraph we present an example of precubical set associated to a trunk.

Nerve of a trunk. We have seen in Example (3.3) that $\square_n$ has a trunk structure and that $\partial^n_{i,\epsilon}$ is a trunk map. Let $\mathcal{T}$ be a trunk, the *nerve of* $\mathcal{T}$ is the precubical set $N(\mathcal{T}) : \square \to \mathbf{Set}$ defined by

$$\begin{aligned} N(\mathcal{T})(\square_n) &:= \mathrm{Hom}_{\mathbf{Trunk}}(\square_n, \mathcal{T}), \\ N(\mathcal{T})(\partial^n_{i,\epsilon}) &:= \mathrm{Hom}_{\mathbf{Trunk}}(-, \mathcal{T})(\partial^n_{i,\epsilon}) = (\partial^n_{i,\epsilon})^*. \end{aligned} \tag{6}$$

In the case of the trunk associated to a rack, the nerve is easy to compute.

Theorem 3.9. *Let X be a rack. There is a bijection $N(X)(\square_n) \overset{\eta_n}{\simeq} X^n$, and under this bijection we have*

$$N(X)(\partial^n_{i,\epsilon})(x_1, \dots, x_n) = \begin{cases} (x_1, \dots, x_{i-1}, x_i \triangleright x_{i+1}, \dots, x_i \triangleright x_n) & \text{if } \epsilon = 0, \\ (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) & \text{if } \epsilon = 1. \end{cases}$$

Lemma 3.10. *If X is a rack, $F \in \mathrm{Hom}_{\mathbf{Trunk}}(\square_n, X)$ and $A \to A \amalg \{b\}$ is an edge in $\square_n$, then*

$$F(A \to A \amalg \{b\}) = \prod_{\substack{1 \le x \le b \\ x \notin A}} F([x-1] \to [x]). \tag{7}$$

where $[x] := \{1, \dots, x\}$.

Proof. (Lemma (3.10)) First recall that in the trunk associated to a rack we have $a \triangleleft b = a$. Suppose $A = \{a_1, \dots, a_p\}$ with $a_1 < \dots < a_p$. We can suppose that $b > a_p$. Indeed, if not then

$$\begin{aligned} F(A \to A \amalg \{b\}) &= F((A \setminus \{a_p\}) \to (A \setminus \{a_p\}) \amalg \{b\}) \triangleleft F((A \setminus \{a_p\}) \amalg \{b\} \to A \amalg \{b\}) \\ &= F((A \setminus \{a_p\}) \to (A \setminus \{a_p\}) \amalg \{b\}) \end{aligned}$$

because

$$\begin{tikzcd} A \arrow[r] & A \amalg \{b\} \\ A \setminus \{a_p\} \arrow[r] \arrow[u] & (A \setminus \{a_p\}) \amalg \{b\} \arrow[u] \end{tikzcd}$$

is a preferred square in $\square_n$. Continuing this reduction until there is no element in A bigger than b make possible this hypothesis.

Now let us show by induction on the cardinal of $\{1, \dots, n\} \setminus A$ the expected equality (7).

- **Initialization** Suppose that the cardinal of $\{1, \dots, n\} \setminus A$ is equal to 1. Because $b > a_p$, necessarily $A = \{1, \dots, n-1\}$ and $b = n$, so the equality (7) is true.
- **Induction:** Suppose the equality true at the rank k. Then for $b_1 = \min\{1 \le x \le b \,|\, x \notin a\}$, by induction hypothesis the following equality holds

$$F(A \amalg \{b_1\} \to A \amalg \{b_1, b\}) = \prod_{\substack{1 \le x \le b \\ x \notin A \amalg \{b_1\}}} F([x-1] \to [x]).$$

Because $b_1 < b$, the square

$$\begin{tikzcd}
A \amalg \{b\} \arrow[r] & A \amalg \{b_1, b\} \\
A \arrow[u] \arrow[r] & A \amalg \{b_1\} \arrow[u]
\end{tikzcd}$$

is preferred in $\square_n$, then

$$F(A \to A \amalg \{b\}) = F(A \to A \amalg \{b_1\}) \rhd F(A \amalg \{b_1\} \to A \amalg \{b_1, b\}).$$

Now using the same reduction as in the begining we have

$$F(A \to A \amalg \{b_1\}) = F(A_{<b_1} \to A_{<b_1} \amalg \{b_1\}),$$

and by definition of b_1 we have $A_{<b_1} = \{1, \dots, b_1 - 1\}$. Thus the expected equality is true.

□

Proof. (Theorem (3.9)) First, let η_n the set map defined from $N(X)(\square_n)$ to X^n by

$$\eta_n : F \mapsto (x_1, \dots, x_n) \tag{8}$$

with $x_{k+1} = F([k] \to [k+1])$ for all $0 \leq k \leq n-1$. By Lemma (3.10) this map is a bijection.

Let $(x_1, \dots, x_n) \in X^n$. Let F be the trunk map in $\mathrm{Hom}_{\mathbf{Trunk}}(\square_n, X)$ corresponding to this n-tuple by the previous bijection. By definition, $N(X)(\partial^n_{i,\epsilon})(x_1, \dots, x_n) = (y_1, \dots, y_{n-1})$ with

$$y_k = (F \circ \partial^n_{i,\epsilon})([k-1] \to [k]).$$

- $\epsilon = 1$: We have

$$\partial^n_{i,1}([k-1] \to [k]) = \begin{cases} [k-1] \amalg \{i\} \to [k] \amalg \{i\} & \text{if } k < i, \\ [k] \to [k+1] & \text{if } k \geq i. \end{cases}$$

Then

$$y_k = \begin{cases} F([k-1] \amalg \{i\} \to [k] \amalg \{i\}) = F([k-1] \to [k]) = x_k & \text{if } k < i, \\ F([k] \to [k+1]) = x_{k+1} & \text{if } k \geq i. \end{cases}$$

- $\epsilon = 0$: We have

$$\partial^n_{i,0}([k-1] \to [k]) = \begin{cases} [k-1] \to [k] & \text{if } k < i, \\ [k] \setminus \{i\} \to [k+1] \setminus \{i\} & \text{if } k \geq i. \end{cases}$$

The square

$$\begin{tikzcd}
{[k+1] \setminus \{i\}} \arrow[r] & {[k+1]} \\
{[k] \setminus \{i\}} \arrow[u] \arrow[r] & {[k]} \arrow[u]
\end{tikzcd}$$

is preferred in $\square_n$, so

$$F([k] \setminus \{i\} \to [k+1] \setminus \{i\}) = F([k] \setminus \{i\} \to [k]) \rhd F([k] \to [k+1]).$$

By Lemma (3.10), this is equal to

$$F([i-1] \to [i]) \rhd F([k] \to [k+1]),$$

and finally

$$y_k = \begin{cases} F([k-1] \to [k]) = x_k & \text{if } k < i, \\ x_i \rhd x_{k+1} & \text{if } k \geq i. \end{cases}$$

$\square$

Let X be a rack and A be an abelian group. By Remark (3.8), the composition $\mathrm{Hom}_{\mathbf{Set}}(-, A) \circ N(X)$ defines a precocubical abelian group. The cochain complex associated to this precocubical abelian group is denoted by $\{C^n(N(X), A), d^n\}_{n\in\mathbb{N}}$.

Corollary 3.11. *Let X be a rack and A be an abelian group (considered as a trivial X-module). There is an isomorphism of cochain complexes*

$$\{CR^n(X, A), d^n_R\}_{n\in\mathbb{N}} \overset{\eta^*}{\simeq} \{C^n(N(X), A), d^n\}_{n\in\mathbb{N}}.$$

4 A graded dendriform algebra structure on $HR^\bullet(X, A)$

Let X be a rack and A be an associative algebra. The goal of this section is to define a graded dendriform algebra structure on $HR^\bullet(X, A)$, the rack cohomology of X with coefficients in A. First we define two graded dendriform products on $\{C^n(N(X), A), d^n\}_{n\in\mathbb{N}}$, the graded module associated to the nerve of a rack X. Then we show that these products are compatible with the differential, and so define a graded dendriform algebra structure on $HR^\bullet(X, A)$.

Products. Let $\sigma \in \mathrm{Sh}_{p_1,p_2}$. In Example (3.5) we defined from σ two trunk maps $\sigma \circ i_{p_1} : \square_{p_1} \to \square_{p_1+p_2}$ and $\sigma \circ i_{p_2} : \square_{p_2} \to \square_{p_1+p_2}$. Let ρ_σ denote the set map from $N(X)(\square_{p_1+p_2})$ to $N(X)(\square_{p_1}) \times N(X)(\square_{p_2})$ defined by

$$\rho_\sigma := ((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2})^*). \tag{9}$$

We define two graded products $\succ$ and $\prec$ on $\{C^n(N(X), A), d^n\}_{n\in\mathbb{N}}$ by

$$f_1 \succ f_2 := \sum_{\sigma\in\mathrm{Sh}^{p_1+p_2}_{p_1,p_2}} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \rho_\sigma, \tag{10}$$

and

$$f_1 \prec f_2 := \sum_{\sigma\in\mathrm{Sh}^{p_1}_{p_1,p_2}} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \rho_\sigma. \tag{11}$$

Dendriform structure. In this paragraph we prove that $\{C^n(N(X), A), \succ, \prec\}_{n\in\mathbb{N}}$ is a graded dendriform algebra. For this we use the bijections

$$\mathrm{Sh}_{p_1,p_2+p_3} \times \mathrm{Sh}_{p_2,p_3} \overset{\alpha}{\simeq} \mathrm{Sh}_{p_1,p_2,p_3} \overset{\beta}{\simeq} \mathrm{Sh}_{p_1+p_2,p_3} \times \mathrm{Sh}_{p_1,p_2}.$$

described in (1).

First, as in Example (3.5), let us define from a shuffle $\sigma \in \mathrm{Sh}_{p_1,p_2,p_3}$ three trunk maps $\sigma \circ i_{p_j} : \square_{p_j} \to \square_{p_1+p_2+p_3}$ where

$$\begin{aligned}
i_{p_1} &:= \partial^{p_1+p_2+p_3}_{p_1+p_2+p_3,0} \circ \cdots \circ \partial^{p_1+1}_{p_1+1,0},\\
i_{p_2} &:= \partial^{p_1+p_2+p_3}_{p_1+p_2+p_3,0} \circ \cdots \circ \partial^{p_1+p_2+1}_{p_1+p_2+1,0} \circ \partial^{p_1+p_2}_{1,1} \circ \cdots \circ \partial^{p_2+1}_{1,1},\\
i_{p_3} &:= \partial^{p_1+p_2+p_3}_{1,1} \circ \cdots \circ \partial^{p_3+1}_{1,1}.
\end{aligned}$$

If $A = \{a_1, \ldots, a_p\} \subseteq \square_{p_j}$, then

$$\begin{aligned}
&(\sigma \circ i_{p_1})(A) = \{\sigma(a_1), \ldots, \sigma(a_p)\},\\
&(\sigma \circ i_{p_2})(A) = \{\sigma(1), \ldots, \sigma(p_1), \sigma(p_1 + a_1), \ldots, \sigma(p_1 + a_p)\},\\
&(\sigma \circ i_{p_3})(A) = \{\sigma(1), \ldots, \sigma(p_1 + p_2), \sigma(p_1 + p_2 + a_1), \ldots, \sigma(p_1 + p_2 + a_p)\}.
\end{aligned}$$

Let ρ_σ the set map from $N(X)(\square_{p_1+p_2+p_3})$ to $N(X)(\square_{p_1}) \times N(X)(\square_{p_2}) \times N(X)(\square_{p_3})$ defined by

$$\rho_\sigma := ((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2})^*, (\sigma \circ i_{p_3})^*).$$

Lemma 4.1. *Let* $\sigma \in \mathrm{Sh}_{p_1,p_2+p_3}$ *(resp.* $\sigma \in \mathrm{Sh}_{p_1+p_2,p_3}$*) and* $\gamma \in \mathrm{Sh}_{p_2,p_3}$ *(resp.* $\gamma \in \mathrm{Sh}_{p_1,p_2}$*). The following equality holds*

$$\rho_{\sigma\circ(1\star\gamma)} = (1 \times \rho_\gamma) \circ \rho_\sigma \text{ (\textit{resp.} } \rho_{\sigma\circ(\gamma\star 1)} = (\rho_\gamma \times 1) \circ \rho_\sigma).$$

Proof. Let $\sigma \in \mathrm{Sh}_{p_1,p_2+p_3}$ and $\gamma \in \mathrm{Sh}_{p_2,p_3}$. Let $F \in N(\mathcal{T})(\square_{p_1+p_2+p_3})$, by definition

$$\rho_{\sigma\circ(1\star\gamma)}(F) = (F \circ (\sigma \circ (1 \star \gamma) \circ i_{p_1}), F \circ (\sigma \circ (1 \star \gamma) \circ i_{p_2}), F \circ (\sigma \circ (1 \star \gamma) \circ i_{p_3}))$$

and

$$((1 \times \rho_\gamma) \circ \rho_\sigma)(F) = (F \circ \sigma \circ i_{p_1}, F \circ \sigma \circ i_{p_2+p_3} \circ \gamma \circ i_{p_2}, F \circ \sigma_{i_{p_2+p_3}} \circ \gamma_{i_{p_3}}).$$

Let $A = \{a_1, \ldots, a_p\} \subseteq \square_{p_1}$.

$$\begin{aligned}
(\sigma \circ (1 \star \gamma) \circ i_{p_1})(A) &= \{(\sigma \circ (1 \star \gamma))(a_1), \ldots, (\sigma \circ (1 \star \gamma))(a_p)\},\\
&= \{\sigma(a_1), \ldots, \sigma(a_p)\},\\
&= (\sigma \circ i_{p_1})(A).
\end{aligned}$$

Let $A = \{a_1, \ldots, a_p\} \subseteq \square_{p_2}$.

$$\begin{aligned}
(\sigma \circ (1 \star \gamma) \circ i_{p_2})(A) &= \{(\sigma \circ (1 \star \gamma))(1), \ldots, (\sigma \circ (1 \star \gamma))(p_1), (\sigma \circ (1 \star \gamma))(p_1 + a_1), \ldots, (\sigma \circ (1 \star \gamma))(p_1 + a_p)\},\\
&= \{\sigma(1), \ldots, \sigma(p_1), \sigma(p_1 + \gamma(a_1)), \ldots, \sigma(p_1 + \gamma(a_p))\},\\
&= (\sigma \circ i_{p_2+p_3} \circ \gamma \circ i_{p_2})(A).
\end{aligned}$$

Let $A = \{a_1, \ldots, a_p\} \subseteq \square_{p_3}$.

$$\begin{aligned}
(\sigma \circ (1 \star \gamma) \circ i_{p_3})(A) &= \{(\sigma \circ (1 \star \gamma))(1), \ldots, (\sigma \circ (1 \star \gamma))(p_1 + p_2), (\sigma \circ (1 \star \gamma))(p_1 + p_2 + a_1), \ldots\\
&\qquad \ldots, (\sigma \circ (1 \star \gamma))(p_1 + p_2 + a_p)\},\\
&= \{\sigma(1), \ldots, \sigma(p_1), \sigma(p_1 + \gamma(1)), \ldots, \sigma(p_1 + \gamma(p_2)), \sigma(p_1 + \gamma(p_2 + a_1)), \ldots, \sigma(p_1 + \gamma(p_2 + a_p))\},\\
&= (\sigma \circ i_{p_2+p_3} \circ \gamma \circ i_{p_3})(A).
\end{aligned}$$

Thus $\rho_{\sigma\circ(1\star\gamma)} = (1 \times \rho_\gamma) \circ \rho_\sigma$. The proof for the other equality is similar. $\square$

Theorem 4.2. *Let X be a rack and A be an associative algebra. The graded module* $\{C^n(N(X), A), \succ, \prec\}_{n\in\mathbb{N}}$ *provided with the products* $\succ$ *and* $\prec$ *is a graded dendriform algebra.*

Proof. Let $p_1, p_2, p_3 \in \mathbb{N}$ and $f_1 \in C^{p_1}(N(X), A), f_2 \in C^{p_2}(N(X), A), f_3 \in C^{p_3}(N(X), A)$. We have to prove the three equalities

1. $f_1 \succ (f_2 \succ f_3) = (f_1 \succ f_2 + f_1 \prec f_2) \succ f_3$,
2. $f_1 \succ (f_2 \prec f_3) = (f_1 \succ f_2) \prec f_3$,

3. $(f_1 \prec f_2) \prec f_3 = f_1 \prec (f_2 \prec f_3 + f_2 \succ f_3)$.

Consider the bijection from $\mathrm{Sh}_{p_1,p_2+p_3} \times \mathrm{Sh}_{p_2,p_3}$ to $\mathrm{Sh}_{p_1+p_2,p_3} \times \mathrm{Sh}_{p_1,p_2}$

$$(\sigma, \gamma) \mapsto (\sigma', \gamma') = (\beta^{-1} \circ \alpha)(\sigma, \gamma).$$

In particular $\epsilon(\sigma)\epsilon(\gamma) = \epsilon(\sigma')\epsilon(\gamma')$.

1. Let $(\sigma, \gamma) \in \mathrm{Sh}^{p_1+p_2+p_3}_{p_1,p_2+p_3} \times \mathrm{Sh}^{p_2+p_3}_{p_2,p_3}$, necessarily $(\sigma', \gamma') \in \mathrm{Sh}^{p_1+p_2+p_3}_{p_1+p_2,p_3} \times \mathrm{Sh}_{p_1,p_2}$. Then, because of cardinality, the map $\beta^{-1} \circ \alpha$ induces a bijection

   $$\mathrm{Sh}^{p_1+p_2+p_3}_{p_1,p_2+p_3} \times \mathrm{Sh}^{p_2+p_3}_{p_2,p_3} \simeq \mathrm{Sh}^{p_1+p_2+p_3}_{p_1+p_2,p_3} \times \mathrm{Sh}_{p_1,p_2}.$$

   Applying the change of variable $\beta^{-1} \circ \alpha$ to the left term in the equality 1. and using the Lemma (4.1), we find the right term in 1..

2. To show this equality, use the bijection $\beta^{-1} \circ \alpha$ restricted to $\mathrm{Sh}^{p_1+p_2+p_3}_{p_1,p_2+p_3} \times \mathrm{Sh}^{p_2}_{p_2,p_3}$. It gives a change of variables

   $$\mathrm{Sh}^{p_1+p_2+p_3}_{p_1,p_2+p_3} \times \mathrm{Sh}^{p_2}_{p_2,p_3} \simeq \mathrm{Sh}^{p_1+p_2}_{p_1+p_2,p_3} \times \mathrm{Sh}^{p_1+p_2}_{p_1,p_2}.$$

3. To show this equality, use the bijection $\beta^{-1} \circ \alpha$ restricted to $\mathrm{Sh}^{p_1}_{p_1,p_2+p_3} \times \mathrm{Sh}_{p_1,p_2}$. It gives a change of variables

   $$\mathrm{Sh}^{p_1}_{p_1,p_2+p_3} \times \mathrm{Sh}_{p_2,p_3} \simeq \mathrm{Sh}^{p_1+p_2}_{p_1+p_2,p_3} \times \mathrm{Sh}^{p_1}_{p_1,p_2}.$$

□

Compatibility of the differential with the products. Let X be a rack and A be an associative algebra. By Theorem (4.2), there is a graded dendriform algebra structure on $C^\bullet(N(X), A)$. In this paragraph we prove that this dendriform structure is compatible with the differential, that is, $\{C^n(N(X), A), d^n, \succ, \prec\}_{n\in\mathbb{N}}$ is a differential graded dendriform algebra.

Theorem 4.3. *Let X be a rack and A be an associative algebra. The cochain complex $\{C^n(N(X), A), d^n\}_{n\in\mathbb{N}}$ provided with the products $\succ$ and $\prec$ is a differential graded dendriform algebra.*

Proof. We have to prove that

$$d^{p_1+1} f_1 \succ f_2 + (-1)^{p_1} f_1 \succ d^{p_2+1} f_2 = d^{p_1+p_2+1}(f_1 \succ f_2) \tag{12}$$

and

$$d^{p_1+1} f_1 \prec f_2 + (-1)^{p_1} f_1 \prec d^{p_2+1} f_2 = d^{p_1+p_2+1}(f_1 \prec f_2) \tag{13}$$

for all $f_1 \in C^{p_1}(N(X), A)$ and $f_2 \in C^{p_2}(N(X), A)$.

To prove these two equalities let us introduce the map ϕ from $\mathbb{S}_{n+1} \times \{1, \dots, n+1\}$ to $\mathbb{S}_n$ defined by

$$\phi(\sigma, i) := (n+1 \dots \sigma(i))\sigma(i \dots n+1).$$

Two important properties of ϕ are the following equalities, satisfied for all $\sigma \in \mathbb{S}_{n+1}, i \in \{1, \dots, p_1 + p_2 + 1\}$ and $\epsilon \in \{0, 1\}$:

$$\partial^{n+1}_{\sigma(i),\epsilon} \circ \phi(\sigma, i) = \sigma \circ \partial^{n+1}_{i,\epsilon}, \tag{14}$$

$$(-1)^{\sigma(i)} \epsilon\big(\phi(\sigma, i)\big) = (-1)^i \epsilon(\sigma). \tag{15}$$

Remark 4.4. The equality (14) is in the category **Gph**.

In the sequel, the restriction to suitable subsets of the map $\psi : \mathbb{S}_{n+1} \times \{1, \dots, n+1\} \to \mathbb{S}_n \times \{1, \dots, n+1\}$ defined by $\psi(\sigma, i) = (\phi(\sigma, i), \sigma(i))$ will provide the good changes of variables needed in order to prove the required equalities.

Compatibility with $\succ$: Let $f_1 \in C^{p_1}(N(X), A)$ and $f_2 \in C^{p_2}(N(X), A)$. The left term in (12) is equal to the sum of four terms (a), (b), (c) and (d) with

$$(\mathrm{a}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}} \sum_{i=1}^{p_1} (-1)^i \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,0})^*, (\sigma \circ i_{p_2})^* \big),$$

$$(\mathrm{b}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}} \sum_{i=1}^{p_2} (-1)^{i+p_1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,0})^* \big),$$

$$(\mathrm{c}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}} \sum_{i=1}^{p_1} (-1)^{i+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,1})^*, (\sigma \circ i_{p_2})^* \big),$$

$$(\mathrm{d}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}} \sum_{i=1}^{p_2} (-1)^{i+p_1+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,1})^* \big).$$

The right term in (12) is equal to the sum of two terms (A) and (B), with

$$(\mathrm{A}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2}_{p_1,p_2}} \sum_{i=1}^{p_1+p_2} (-1)^i \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\partial^{p_1+p_2+1}_{i,0} \circ \sigma \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{i,0} \circ \sigma \circ i_{p_2})^* \big),$$

$$(\mathrm{B}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2}_{p_1,p_2}} \sum_{i=1}^{p_1+p_2} (-1)^{i+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\partial^{p_1+p_2+1}_{i,1} \circ \sigma \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{i,1} \circ \sigma \circ i_{p_2})^* \big).$$

In order to prove (12), we are going to show that

$$(\mathrm{a}) + (\mathrm{c}) + (\mathrm{d}) = (\mathrm{B}) \ \text{ and } \ (\mathrm{b}) = (\mathrm{A}).$$

•(b) = (A) : This equality is proved using the change of variables

$$\psi \circ t_{p_1} : \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1} \times \{1, \dots, p_2\} \to \mathrm{Sh}^{p_1+p_2}_{p_1,p_2} \times \{1, \dots, p_1+p_2\}$$

with $t_{p_1}(i) = p_1 + i$. Using this change of variables, we have to prove for all $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}$ and $i \in \{1, \dots, p_2\}$:

1. $\big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,0})^*\big) = \big((\partial^{p_1+p_2+1}_{\sigma(p_1+i),0} \circ \phi(\sigma, p_1+i) \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{\sigma(p_1+i),0} \circ \phi(\sigma, p_1+i) \circ i_{p_2})^*\big)$,
2. $(-1)^{\sigma(p_1+i)} \epsilon\big(\phi(\sigma, p_1+i)\big) = (-1)^{p_1+i} \epsilon(\sigma)$.

1. Let $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}$ and $i \in \{1, \dots, p_2\}$. We want to prove for all $F \in \mathrm{Hom}_{\mathbf{Trunk}}(\square_{p_1+p_2+1}, X)$

$$\begin{cases} F \circ \sigma \circ i_{p_1} = F \circ \partial^{p_1+p_2+1}_{\sigma(p_1+i),0} \circ \phi(\sigma, p_1+i) \circ i_{p_1}, \\ F \circ \sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,0} = F \circ \partial^{p_1+p_2+1}_{\sigma(p_1+i),0} \circ \phi(\sigma, p_1+i) \circ i_{p_2}. \end{cases}$$

By Theorem (3.9), this is equivalent to prove

$$\left\{\begin{array}{ll} (F\circ\sigma\circ i_{p_1})([k-1]\to[k]) = (F\circ\partial^{p_1+p_2+1}_{\sigma(p_1+i),0}\circ\phi(\sigma,p_1+i)\circ i_{p_1})([k-1]\to[k]) & 1\le k\le p_1, \\ (F\circ\sigma\circ i_{p_2+1}\circ\partial^{p_2+1}_{i,0})([k-1]\to[k]) = (F\circ\partial^{p_1+p_2+1}_{\sigma(p_1+i),0}\circ\phi(\sigma,p_1+i)\circ i_{p_2})([k-1]\to[k]) & 1\le k\le p_2. \end{array}\right.$$

Let $0\le k\le p_1$, $(\sigma\circ i_{p_1})([k]) = \sigma([k])$ and $(\partial^{p_1+p_2+1}_{\sigma(p_1+i),0}\circ\phi(\sigma,p_1+i)\circ i_{p_1})([k]) \overset{(14)}{=} (\sigma\circ\partial^{p_1+p_2+1}_{p_1+i,0}\circ i_{p_1})([k]) = \sigma([k])$, so the first equality.
Let $0\le k\le p_2$,

$$\begin{aligned} (\sigma\circ i_{p_2+1}\circ\partial^{p_2+1}_{i,0})([k]) &= \left\{\begin{array}{ll} (\sigma\circ i_{p_2+1})([k]) & \text{if } k<i, \\ (\sigma\circ i_{p_2+1})([k+1]\setminus\{i\}) & \text{if } k\ge i, \end{array}\right. \\ &= \left\{\begin{array}{ll} \sigma([p_1+k]) & \text{if } k<i, \\ \sigma([p_1+k+1]\setminus\{p_1+i\}) & \text{if } k\ge i, \end{array}\right. \end{aligned}$$

and

$$\begin{aligned} \partial^{p_1+p_2+1}_{\sigma(p_1+i),0}\circ\phi(\sigma,p_1+i)\circ i_{p_2})([k]) &= \partial^{p_1+p_2+1}_{\sigma(p_1+i),0}\circ\phi(\sigma,p_1+i))([p_1+k]) \\ &\overset{(14)}{=} (\sigma\circ\partial^{p_1+p_2+1}_{p_1+i,0})([p_1+k]) \\ &= \left\{\begin{array}{ll} \sigma([p_1+k]) & \text{if } k<i, \\ \sigma([p_1+k+1]\setminus\{p_1+i\}) & \text{if } k\ge i, \end{array}\right. \end{aligned}$$

so the second equality.

2. Deduced from (15).

•(a) + (c) + (d) = (B) : To prove this equality we need to decompose (d) into two sums. Let us denote by $I^{\ge}$ and $I^{<}$ the subsets of $\mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}\times\{1,\dots,p_2\}$ defined by

$$\begin{aligned} I^{\ge} &:= \{(\sigma,i)\in\mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}\times\{1,\dots,p_2\}\,|\,\sigma(p_1+i)\ge p_1+i\} \\ &= \{(\sigma,i)\in\mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}\times\{1,\dots,p_2\}\,|\,\sigma(p_1+i)= p_1+i\} \end{aligned}$$

and

$$I^{<} := (\mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}\times\{1,\dots,p_2\})\setminus I^{\ge}.$$

Then we are going to prove that

$$(\mathrm{a}) + \sum_{I^{<}} = 0 \ \text{ and } \ (\mathrm{c}) + \sum_{I^{\ge}} = (\mathrm{B}).$$

(a) + $\sum_{I^{<}} = 0$: To prove this equality we use the change of variables

$$\theta : \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}\times\{1,\dots,p_1\}\to I^{<}$$

defined by $\theta(\sigma,i) = (\rho(\sigma,i), 1+\sigma(i)-i)$ with $\rho(\sigma,i) = \sigma\circ(i\dots p_1+1+\sigma(i)-i)$. Using this change of variables, we have to prove for all $\sigma\in\mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}$ and $i\in\{1,\dots,p_1\}$:

1. $\big((\sigma\circ i_{p_1+1}\circ\partial^{p_1+1}_{i,0})^*, (\sigma\circ i_{p_2})^*\big) = \big((\rho(\sigma,i)\circ i_{p_1})^*, (\rho(\sigma,i)\circ i_{p_2+1}\circ\partial^{p_2+1}_{1+\sigma(i)-i,1})^*\big)$,
2. $(-1)^{p_1+1+\sigma(i)-i}\epsilon\big(\rho(\sigma,i)\big) = (-1)^i\epsilon(\sigma)$.

1. Let $\sigma \in \mathrm{Sh}_{p_1+1,p_2}^{p_1+p_2+1}$ and $i \in \{1,\dots,p_1\}$. Using Theorem (3.9), we are going to prove, for all $F \in \mathrm{Hom}_{\mathbf{Trunk}}(\square_{p_1+p_2+1}, X)$, the following equalities :

$$\left\{\begin{array}{ll} (F\circ\sigma\circ i_{p_1+1}\circ\partial_{i,0}^{p_1+1})([k-1]\to[k]) = (F\circ\rho(\sigma,i)\circ i_{p_1})([k-1]\to[k]) & \text{if } 1\le k\le p_1, \\ (F\circ\sigma\circ i_{p_2})([k-1]\to[k]) = (F\circ\rho(\sigma,i)\circ i_{p_2+1}\circ\partial_{1+\sigma(i)-i,1}^{p_2+1})([k-1]\to[k]) & \text{if } 1\le k\le p_2. \end{array}\right.$$

Let $0\le k\le p_1$,

$$(\sigma\circ i_{p_1+1}\circ\partial_{i,0}^{p_1+1})([k]) = \left\{\begin{array}{ll} \sigma([k]) & \text{if } k<i, \\ \sigma([k+1]\setminus\{i\}) & \text{if } k\ge i. \end{array}\right.$$

and

$$\begin{aligned}(\rho(\sigma,i)\circ i_{p_1})([k]) &= \big(\sigma\circ(i\dots p_1+1+\sigma(i)-i)\big)([k]),\\ &= \left\{\begin{array}{ll} \sigma([k]) & \text{if } k<i, \\ \sigma([k+1]\setminus\{i\}) & \text{if } k\ge i, \end{array}\right.\end{aligned}$$

(remark that $\sigma(i)-i\ge 0$ for $i\in\{1,\dots,p_1\}$), so the first equality.
Let $0\le k\le p_2$, we have $(\sigma\circ i_{p_2})([k]) = \sigma([p_1+1+k])$ and

$$\begin{aligned}(\rho(\sigma,i)\circ i_{p_2+1}\circ\partial_{1+\sigma(i)-i,1}^{p_2+1})([k]) &= \left\{\begin{array}{ll} (\rho(\sigma,i)\circ i_{p_2+1})([k]\amalg\{1+\sigma(i)-i\} & \text{if } k<1+\sigma(i)-i, \\ (\rho(\sigma,i)\circ i_{p_2+1})([k+1]) & \text{if } k\ge 1+\sigma(i)-i, \end{array}\right.\\ &= \left\{\begin{array}{ll} \rho(\sigma,i)([p_1+k]\amalg\{p_1+1+\sigma(i)-i\}) & \text{if } k<1+\sigma(i)-i, \\ \rho(\sigma,i)([p_1+k+1]) & \text{if } k\ge 1+\sigma(i)-i, \end{array}\right.\\ &= \left\{\begin{array}{ll} \sigma([p_1+k+1])\amalg\{\sigma(i)\} & \text{if } k<1+\sigma(i)-i, \\ \sigma([p_1+k+1]) & \text{if } k\ge 1+\sigma(i)-i, \end{array}\right.\\ &= \sigma([p_1+k+1]),\end{aligned}$$

so the second equality.

2. Clear by definition of $\rho(\sigma,i)$.

(c) $+\displaystyle\sum_{I^{\geq}}$ = (B) : To prove this equality we need to decompose (B) into two sums, one equals to $\sum_{I^{\geq}}$ and the other equals to (c). Let us denote by $J^{\geq}$ and $J^{<}$ the subsets of $\mathrm{Sh}_{p_1,p_2}^{p_1+p_2}\times\{p_1+1,\dots,p_1+p_2\}$ defined by

$$J^{\geq} := \{(\sigma,i)\in\mathrm{Sh}_{p_1,p_2}^{p_1+p_2}\times\{p_1+1,\dots,p_1+p_2\}\,|\,\sigma(i)\ge i\}$$

and

$$J^{<} := (\mathrm{Sh}_{p_1,p_2}^{p_1+p_2}\times\{p_1+1,\dots,p_1+p_2\})\setminus J^{\geq}.$$

Then we are going to prove that

$$\sum_{I^{\geq}} = \sum_{J^{\geq}} \quad\text{and}\quad \text{(c)} = \sum_{J^{<}}.$$

$\sum_{I^{\geq}} = \sum_{J^{\geq}}$: This equality is proved using the change of variables

$$\psi\circ t_{p_1} : I^{\geq}\to J^{\geq}.$$

In this case we have $(\psi \circ t_{p_1})(\sigma, i) = (\sigma, p_1 + i)$. Using this change of variables, we have to prove for all $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}$ and $i \in \{1, \dots, p_2\}$:

$$\big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,1})^*\big) = \big((\partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_2})^*\big).$$

Let $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1,p_2+1}$ and $i \in \{1, \dots, p_2\}$. Using Theorem (3.9), we are going to prove, for all $F \in \mathrm{Hom}_{\mathbf{Trunk}}(\square_{p_1+p_2+1}, X)$, the following equalities :

$$\left\{\begin{array}{ll} (F \circ \sigma \circ i_{p_1})([k-1] \to [k]) = (F \circ \partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_1})([k-1] \to [k]) & \text{if } 1 \leq k \leq p_1, \\ (F \circ \sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,1})([k-1] \to [k]) = (F \circ \partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_2})([k-1] \to [k]) & \text{if } 1 \leq k \leq p_2. \end{array}\right.$$

Let $0 \leq k \leq p_1$, $(\sigma \circ i_{p_1})([k]) = \sigma([k])$ and $(\partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_1})([k]) \overset{(14)}{=} (\sigma \circ \partial^{p_1+p_2+1}_{p_1+i,1} \circ i_{p_1})([k]) = \sigma([k]) \amalg \{\sigma(p_1+i)\} = \sigma([k]) \amalg \{p_1 + i\}$. Hence the first equality is equivalent to

$$F(\sigma([k-1]) \to \sigma([k])) = F\big((\sigma([k-1]) \amalg \{p_1+i\}) \to (\sigma([k]) \amalg \{p_1+i\})\big).$$

Because $\sigma \in I^{\geq}$, necessarily $\sigma(k) < p_1 + i$, so the square

$$\begin{tikzcd}
\sigma([k-1]) \amalg \{p_1+i\} \arrow[r] & \sigma([k]) \amalg \{p_1+i\} \\
\sigma([k-1]) \arrow[r] \arrow[u] & \sigma([k]) \arrow[u]
\end{tikzcd}$$

is a preferred square in $\square_{p_1+p_2+1}$, and the expected equality si proved.
Let $0 \leq k \leq p_2$, we have

$$(\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,1})([k]) = \left\{\begin{array}{ll} \sigma([p_1+k]) \amalg \{p_1+i\} & \text{if } k < i, \\ \sigma([p_1+k+1]) & \text{if } k \geq i, \end{array}\right.$$

and

$$\begin{aligned}(\partial^{p_1+p_2+1}_{p_1+i,1} \circ \sigma \circ i_{p_2})([k]) &\overset{(14)}{=} (\sigma \circ \partial^{p_1+p_2+1}_{p_1+i,1} \circ i_{p_2})([k]), \\ &= \left\{\begin{array}{ll} \sigma([p_1+k]) \amalg \{p_1+i\} & \text{if } k < i, \\ \sigma([p_1+k+1]) & \text{if } k \geq i, \end{array}\right.\end{aligned}$$

so the second equality.

(c) $= \sum_{J^<}$: The following equality is proved using the change of variables

$$\psi : \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2} \times \{1, \dots, p_1\} \to J^<.$$

Using this change of variables, we have to prove for all $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}$ and $i \in \{1, \dots, p_1\}$:

1. $\big((\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,1})^*, (\sigma \circ i_{p_2})^*\big) = \big((\partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma,i) \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma,i) \circ i_{p_2})^*\big)$,
2. $(-1)^{i+1}\epsilon(\sigma) = (-1)^{\sigma(i)+1}\epsilon(\phi(\sigma,i))$.

1. Let $\sigma \in \mathrm{Sh}^{p_1+p_2+1}_{p_1+1,p_2}$ and $i \in \{1, \dots, p_1\}$. Using Theorem (3.9), we are going to prove, for all $F \in \mathrm{Hom}_{\mathbf{Trunk}}(\square_{p_1+p_2+1}, X)$, the following equalities :

$$\left\{\begin{array}{ll} (F \circ \sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,1})([k-1] \to [k]) = (F \circ \partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma,i) \circ i_{p_1})([k-1] \to [k]) & \text{if } 1 \leq k \leq p_1, \\ (F \circ \sigma \circ i_{p_2})([k-1] \to [k]) = (F \circ \partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma,i) \circ i_{p_2})([k-1] \to [k]) & \text{if } 1 \leq k \leq p_2. \end{array}\right.$$

Let $0 \leq k \leq p_1$, we have

$$(\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,1})([k]) = \begin{cases} \sigma([k]) \amalg \{\sigma(i)\} & \text{if } k < i, \\ \sigma([k+1]) & \text{if } k \geq i, \end{cases}$$

and

$$\begin{aligned}(\partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma, i) \circ i_{p_1})([k]) &\overset{(14)}{=} (\sigma \circ \partial^{p_1+p_2+1}_{i,1} \circ i_{p_1})([k]), \\ &= \begin{cases} \sigma([k]) \amalg \{\sigma(i)\} & \text{if } k < i, \\ \sigma([k+1]) & \text{if } k \geq i, \end{cases}\end{aligned}$$

so the first equality.
Let $0 \leq k \leq p_2$, we have $(\sigma \circ i_{p_2})([k]) = \sigma([p_1 + 1 + k])$ and $(\partial^{p_1+p_2+1}_{\sigma(i),1} \circ \phi(\sigma, i) \circ i_{p_2})([k]) \overset{(14)}{=} (\sigma \circ \partial^{p_1+p_2+1}_{i,1} \circ i_{p_2})([k]) = (\sigma \circ \partial^{p_1+p_2+1}_{i,1})([p_1 + k]) = \sigma([p_1 + 1 + k])$, so the second equality.

2. Deduced from (15).

Compatibility with $\prec$: The proof is essentially the same as before. Let $f_1 \in C^{p_1}(N(X), A)$ and $f_2 \in C^{p_2}(N(X), A)$. The left term in (13) is equal to the sum of four terms (a), (b), (c) and (d) with

$$\text{(a)} = \sum_{\sigma \in \mathrm{Sh}^{p_1+1}_{p_1+1,p_2}} \sum_{i=1}^{p_1} (-1)^i \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,0})^*, (\sigma \circ i_{p_2})^*\big),$$

$$\text{(b)} = \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2+1}} \sum_{i=1}^{p_2} (-1)^{i+p_1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,0})^*\big),$$

$$\text{(c)} = \sum_{\sigma \in \mathrm{Sh}^{p_1+1}_{p_1+1,p_2}} \sum_{i=1}^{p_1} (-1)^{i+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1+1} \circ \partial^{p_1+1}_{i,1})^*, (\sigma \circ i_{p_2})^*\big),$$

$$\text{(d)} = \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2+1}} \sum_{i=1}^{p_2} (-1)^{i+p_1+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\sigma \circ i_{p_1})^*, (\sigma \circ i_{p_2+1} \circ \partial^{p_2+1}_{i,1})^*\big).$$

The right term in (13) is equal to the sum of two terms (A) and (B), with

$$\text{(A)} = \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2}} \sum_{i=1}^{p_1+p_2} (-1)^i \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\partial^{p_1+p_2+1}_{i,0} \circ \sigma \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{i,0} \circ \sigma i_{p_2})^*\big),$$

$$\text{(B)} = \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2}} \sum_{i=1}^{p_1+p_2} (-1)^{i+1} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ \big((\partial^{p_1+p_2+1}_{i,1} \circ \sigma \circ i_{p_1})^*, (\partial^{p_1+p_2+1}_{i,1} \circ \sigma \circ i_{p_2})^*\big).$$

In order to prove (13), we are going to show that

$$\text{(a)} + \text{(b)} + \text{(d)} = \text{(A)} \;\text{ and }\; \text{(c)} = \text{(B)}.$$

•(c) = (B) : The following equality is proved using the change of variables

$$\psi : \mathrm{Sh}^{p_1+1}_{p_1+1,p_2} \times \{1, \dots, p_1\} \to \mathrm{Sh}^{p_1}_{p_1,p_2} \times \{1, \dots, p_1 + p_2\}.$$

$\bullet$(a) + (b) + (d) = (A) : To prove this equality we need to decompose (a) into two sums. Let us denote by $I^{\geq}$ and $I^{<}$ the subsets of $\mathrm{Sh}^{p_1+1}_{p_1+1,p_2} \times \{1,\ldots,p_1\}$ defined by

$$\begin{aligned} I^{\geq} &:= \{(\sigma,i) \in \mathrm{Sh}^{p_1+1}_{p_1+1,p_2} \times \{1,\ldots,p_1\} \,|\, \sigma(i) \geq p_2 + i\}, \\ &= \{(\sigma,i) \in \mathrm{Sh}^{p_1+1}_{p_1+1,p_2} \times \{1,\ldots,p_1\} \,|\, \sigma(i) = p_2 + i\}. \end{aligned}$$

and

$$I^{<} := (\mathrm{Sh}^{p_1+1}_{p_1+1,p_2} \times \{1,\ldots,p_1\}) \setminus I^{<}.$$

Then we are going to show that

$$(\mathrm{d}) + \sum_{I^{<}} = 0 \ \text{ and } \ (\mathrm{b}) + \sum_{I^{\geq}} = (\mathrm{A}).$$

(d) + $\sum_{I^{<}} = 0$: The following equality is proved using the change of variables

$$\theta : \mathrm{Sh}^{p_1}_{p_1,p_2+1} \times \{1,\ldots,p_2\} \to I^{<}$$

defined by $\theta(\sigma,i) = (\rho(\sigma,i), \sigma(p_1+i)-i+1)$ with $\rho(\sigma,i) = \sigma \circ (p_1+i\; p_1+i-1 \ldots \gamma(p_1+i)-i+1)$.

(b) + $\sum\limits_{I^{\geq}}$ = (A) : To prove this equality we need to decompose (A) into two sums, one equals to $\sum_{I^{\geq}}$ and the other equals to (b). Let us denote by $J^{\geq}$ and $J^{<}$ the subsets of $\mathrm{Sh}^{p_1}_{p_1,p_2} \times \{1,\ldots,p_1\}$ defined by

$$J^{\geq} := \{(\sigma,i) \in \mathrm{Sh}^{p_1}_{p_1,p_2} \times \{p_2+1,\ldots,p_1+p_2\} \,|\, \sigma(i-p_2) \geq i\}$$

and

$$J^{<} := (\mathrm{Sh}^{p_1}_{p_1,p_2} \times \{1,\ldots,p_1\}) \setminus J^{\geq}.$$

Then we are going to show that

$$\sum_{I^{\geq}} = \sum_{I^{\geq}} \ \text{ and } \ (\mathrm{b}) = \sum_{J^{<}}.$$

$\sum_{I^{\geq}} = \sum_{J^{\geq}}$: The following equality is proved using the change of variables

$$\psi : I^{\geq} \to J^{\geq}.$$

(b) = $\sum_{J^{<}}$: The following equality is proved using the change of variables

$$\psi \circ t_{p_1} : \mathrm{Sh}^{p_1}_{p_1,p_2+1} \times \{1,\ldots,p_2\} \to J^{<}.$$

$\square$

Explicit formulas for $\succ$ and $\prec$. Using the cochain complex isomorphism η^* between $\{CR^n(X,A), d^n_R\}_{n\in\mathbb{N}}$ and $\{C^n(N(X),A), d^n\}_{n\in\mathbb{N}}$ (Corollary (3.11)), the differential graded dendriform algebra structure on $\{CR^n(X,A), d^n_R\}_{n\in\mathbb{N}}$ is defined by the formulas :

$$f_1 \succ f_2 := \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2}_{p_1,p_2}} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ (\eta_{p_1} \times \eta_{p_2}) \circ \rho_\sigma \circ \eta^{-1}_{p_1+p_2},$$

and

$$f_1 \prec f_2 := \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2}} \epsilon(\sigma)\, \mu_A \circ (f_1 \times f_2) \circ (\eta_{p_1} \times \eta_{p_2}) \circ \rho_\sigma \circ \eta^{-1}_{p_1+p_2}.$$

Then to have formulas for $\succ$ and $\prec$ on $\{CR^n(X,A), d^n_R\}$ we have to compute

$$(\eta_{p_1} \times \eta_{p_2}) \circ \rho_\sigma \circ \eta^{-1}_{p_1+p_2} : X^{p_1+p_2} \to X^{p_1} \times X^{p_2}.$$

Let $(x_1, \dots, x_{p_1+p_2}) \in X^{p_1+p_2}$, by (8) and (9)

$$\big((\eta_{p_1} \times \eta_{p_2}) \circ \rho_\sigma \circ \eta^{-1}_{p_1+p_2}\big)(x_1, \dots, x_{p_1+p_2}) = \big((y_1, \dots, y_{p_1}), (z_1, \dots, z_{p_2})\big),$$

where

$$y_k = (F \circ \sigma \circ i_{p_1})([k-1] \to [k]) \text{ and } z_k = (F \circ \sigma \circ i_{p_2})([k-1] \to [k]).$$

Thus, using Lemma (3.10)

$$\begin{aligned} y_k &= (F \circ \sigma)([k-1] \to [k]), \\ &= F\big(\sigma([k-1]) \to \sigma([k])\big), \\ &= F\big(\sigma([k-1]) \to \sigma([k-1] \amalg \{\sigma(k)\})\big), \\ &= \prod_{\substack{1 \leq x \leq \sigma(k) \\ x \notin \sigma([k-1])}} F([x-1] \to [x]), \\ &= x_{i_1} \triangleright \cdots \triangleright x_{i_j} \triangleright x_{\sigma(k)}, \end{aligned}$$

with $x_{i_1} < \cdots < x_{i_j} < \sigma(k)$ and $i_l \in \{\sigma(p_1+1), \dots, \sigma(p_1+p_2)\}$. In the same way

$$\begin{aligned} z_k &= (F \circ \sigma)([p_1+k-1] \to [p_1+k]), \\ &= F\big(\sigma([p_1+k-1]) \to \sigma([p_1+k])\big), \\ &= F\big(\sigma([p_1+k-1]) \to \sigma([p_1+k-1] \amalg \{\sigma(p_1+k)\})\big), \\ &= \prod_{\substack{1 \leq x \leq \sigma(p_1+k) \\ x \notin \sigma([p_1+k-1])}} F([x-1] \to [x]), \\ &= x_{\sigma(p_1+k)}. \end{aligned}$$

Finally we have proved the following theorem.

Theorem 4.5. *Let X be a rack and A be an associative algebra. The cochain complex $\{CR^n(X,A), d^n_R\}_{n \in \mathbb{N}}$ provided with the products $\succ$ and $\prec$ defined by*

$$(f_1 \succ f_2)(x_1, \dots, x_{p_1+p_2}) = \sum_{\sigma \in \mathrm{Sh}^{p_1+p_2}_{p_1,p_2}} \epsilon(\sigma)\, f_1(y_1, \dots, y_{p_1}) f_2(z_1, \dots, z_{p_2}),$$

$$(f_1 \prec f_2)(x_1, \dots, x_{p_1+p_2}) = \sum_{\sigma \in \mathrm{Sh}^{p_1}_{p_1,p_2}} \epsilon(\sigma)\, f_1(y_1, \dots, y_{p_1}) f_2(z_1, \dots, z_{p_2}),$$

where

$$\begin{cases} y_k = x_{i_1} \triangleright \cdots \triangleright x_{i_j} \triangleright x_{\sigma(k)}, \\ z_k = x_{\sigma(p_1+k)}, \end{cases}$$

with $x_{i_1} < \cdots < x_{i_j} < \sigma(k)$ and $i_l \in \{\sigma(p_1+1), \dots, \sigma(p_1+p_2)\}$, is a differential graded dendriform algebra. In particular $HR^\bullet(X,A)$ is provided with a graded dendriform algebra structure.

Remark 4.6. If the rack is pointed, then remark that these formulas are well defined on the cochain complex $\{CR^n_p(X,A), d^n_R\}_{n \in \mathbb{N}}$. Notice that these formulas hold also in the case of a shelf (resp. pointed shelf).

5 A graded associative algebra morphism from $H^\bullet(G,A)$ to $HR^\bullet(\mathrm{Conj}(G),A)$

Let G be group and A be an associative group. The cochain complex $\{C^n(G,A), d^n_G, \cup\}$ calculating the group cohomology with coefficient in A is provided with a differential graded associative algebra structure given by the cup product $\cup$. Moreover, considering the rack Conj(G) associated to G (Example (2.1)), we have shown (Theorem (4.5)) that the cochain complex $\{CR^n(\mathrm{Conj(G)},A), d^n_R, \star\}$, with $\star = \succ + \prec$, is a differential graded associative algebra. In this section we define a differential graded associative algebra morphism from $\{C^n(G,A), d^n_G, \cup\}$ to $\{CR^n(\mathrm{Conj(G)},A), d^n_R, \star\}$.

Group cohomology. Let G be a group and A be a left G-module. The cochain complex $\{C^n(G,A), d^n_G\}_{n\in\mathbb{N}}$ is defined by

$$\begin{aligned} C^n(G,A) &:= \mathrm{Hom}_{\mathbf{Set}}(G^n, A), \\ d^{n+1}_G &:= \sum_{i=0}^{n+1} (-1)^i d^{n+1}_i f, \end{aligned}$$

where

$$d^{n+1}_i f(x_1,\ldots,x_{n+1}) := \begin{cases} x_1 \cdot f(x_2,\ldots,x_{n+1}) & \text{if } i=0, \\ f(x_1,\ldots,x_i x_{i+1},\ldots,x_{n+1}) & \text{if } 1 \le i \le n, \\ f(x_1,\ldots,x_n) & \text{if } i = n+1. \end{cases}$$

The family of maps $\{d^n_i\}$ satisfied the *simplicial identities*, that is, for all $0 \le i < j \le n+1$, we have the identities :

$$d^{n+1}_i \circ d^n_{j-1} = d^{n+1}_j \circ d^n_i,$$

so d_G is a differential.

The cohomology associated to this cochain complex is called the *group cohomology of* G *with coefficients in* A and denoted $H^\bullet(G,A)$.

Cup product on group cohomology. Let G be a group and A be an associative algebra with product denoted by μ_A. Let $p_1, p_2 \in \mathbb{N}$, let ρ denote the set map from $G^{p_1+p_2}$ to $G^{p_1} \times G^{p_2}$ defined by

$$\rho(x_1,\ldots,x_{p_1+p_2}) := \big((x_1,\ldots,x_{p_1}),(x_{p_1+1},\ldots,x_{p_1+p_2})\big).$$

A graded product $\cup$, called *cup product*, is defined on $\{C^n(G,A), d^n_G\}_{n\in\mathbb{N}}$ by the formula

$$f_1 \cup f_2 := \mu_A \circ (f_1 \times f_2) \circ \rho.$$

Theorem 5.1. *Let G be a group and A be an associative algebra (considered as a trivial G-module). The cochain complex $\{C^n(G,A), d^n_G\}_{n\in\mathbb{N}}$ provided with the cup product $\cup$ is a differential graded associative algebra structure.*

Equivalent definitions of the group cohomology and the cup product. Let $G_{\mathbf{Cat}}$ denote the category canonically associated to a group G (Example (3.1)). Let $\mathrm{B}(G)$ denote its *presimplicial nerve of* $G_{\mathbf{Cat}}$, that is, the functor from Δ to **Set** defined by

$$\begin{aligned} \mathrm{B}(G)(\Delta_n) &:= \mathrm{Hom}_{\mathbf{Cat}}(\Delta_n, G_{\mathbf{Cat}}), \\ \mathrm{B}(G)(\partial^n_i) &:= \mathrm{Hom}_{\mathbf{Cat}}(-, G_{\mathbf{Cat}})(\partial^n_i) = (\partial^n_i)^*. \end{aligned}$$

Proposition 5.2. *Let G be a group. There is a bijection $\mathrm{B}(G)(\Delta_n) \overset{\lambda_n}{\simeq} G^n$, and under this bijection we have*

$$(\partial_i^n)^*(x_1,\dots,x_n) = \begin{cases} (x_2,\dots,x_{n+1}) & \text{if } i=0, \\ (x_1,\dots,x_i x_{i+1},\dots,x_{n+1}) & \text{if } 1 \le i \le n, \\ (x_1,\dots,x_n) & \text{if } i = n+1. \end{cases}$$

Let A be a left G-module. The composition $\mathrm{Hom}_{\mathbf{Set}}(-,A) \circ \mathrm{B}(G)$ defines a precosimplicial abelian group . Denote by $\{C_\Delta^n(G,A), d_\Delta^n\}_{n\in\mathbb{N}}$ the cochain complex associated to this precosimplicial abelian group. The cohomology associated to this cochain complex is denoted $H_\Delta^\bullet(G,A)$.

Corollary 5.3. *Let G be a group and A be an abelian group (considered as a trivial left G-module). There is an isomorphism of cochain complexes*

$$\{C^n(G,A), d_G^n\}_{n\in\mathbb{N}} \overset{\lambda^*}{\simeq} \{C_\Delta^n(G,A), d_\Delta^n\}_{n\in\mathbb{N}}.$$

Let $p_1, p_2 \in \mathbb{N}$. We define two functors $j_{p_1} : \Delta_{p_1} \to \Delta_{p_1+p_2}$ and $j_{p_2} : \Delta_{p_2} \to \Delta_{p_1+p_2}$ by $j_{p_1}(k) = k$ and $j_{p_2}(k) = p_1 + k$. Let ρ denote the set map from $\mathrm{B}(G)(\Delta_{p_1+p_2})$ to $\mathrm{B}(G)(\Delta_{p_1}) \times \mathrm{B}(G)(\Delta_{p_2})$ defined by

$$\rho := (j_{p_1}^*, j_{p_2}^*).$$

A graded product $\cup_\Delta$ on $\{C_\Delta^n(G,A), d_\Delta^n\}_{n\in\mathbb{N}}$ is defined by

$$f_1 \cup_\Delta f_2 := \mu_A \circ (f_1 \times f_2) \circ \rho.$$

Proposition 5.4. *Let G be a group and A be an associative algebra (considered as a trivial G-module). The cochain complex $\{C_\Delta^n(G,A), d_\Delta^n\}_{n\in\mathbb{N}}$ provided with the product $\cup_\Delta$ is a differential graded associative algebra.*

Proposition 5.5. *Let G be a group and A be an associative algebra (considered as a trivial G-module). The cochain complex isomorphism between $\{C^n(G,A), d_G^n\}_{n\in\mathbb{N}}$ and $\{C_\Delta^n(G,A), d_\Delta^n\}_{n\in\mathbb{N}}$ given in Corollary* (5.3) *is a differential graded associative algebra isomorphism.*

Cubical cohomology of groups. We have seen in Example (3.1) and (3.2) that $\Box_n$ has a category structure and that $\partial_{i,\epsilon}^n$ is a functor. Hence, there is a precubical set $N(\mathcal{C}) : \Box \to \mathbf{Set}$ associated to any category $\mathcal{C}$, called *the cubical nerve of* $\mathcal{C}$, and defined by

$$\begin{aligned} N(\mathcal{C})(\Box_n) &:= \mathrm{Hom}_{\mathbf{Cat}}(\Box_n, \mathcal{C}), \\ N(\mathcal{C})(\partial_{i,\epsilon}^n) &:= \mathrm{Hom}_{\mathbf{Cat}}(-,\mathcal{C})(\partial_{i,\epsilon}^n) = (\partial_{i,\epsilon}^n)^*. \end{aligned}$$

Let G be a group and A be a left G-module. The composition $\mathrm{Hom}_{\mathbf{Set}}(-,A) \circ N(G_{\mathbf{Cat}})$ defines a precocubical abelian group. Denote by $\{C_\Box^n(G,A), d_\Box^n\}_{n\in\mathbb{N}}$ the cochain complex associated to this precocubical abelian group. The cohomology associated to this cochain complex is called the *cubical cohomology of G with coefficient in A* and denoted $H_\Box^\bullet(G,A)$.

The morphism $H^\bullet(G,A) \overset{[S^\bullet]}{\to} HR^\bullet(G,A)$. Define a graded abelian group morphism $S^\bullet$ as the following composition :

$$\begin{tikzcd} C^\bullet(G,A) \arrow[rrrr, dotted, "S^\bullet"] \arrow[d, "\lambda^\bullet"'] & & & & CR^\bullet(\mathrm{Conj}(G),A) \\ C_\Delta^\bullet(G,A) \arrow[r, "\Sigma^\bullet"] & C_\Box^\bullet(G,A) \arrow[r, "T^\bullet"] & C^\bullet(N(\mathrm{Tr}(G_{\mathbf{Cat}})),A) \arrow[r, "I^\bullet"] & C^\bullet(N(\mathrm{Conj}(G)),A) \arrow[r, phantom] & \\ \end{tikzcd} \tag{16}$$

where

- $I^n = \big(\mathrm{Hom}_{\mathbf{Set}}(-,A)\circ\mathrm{Hom}_{\mathbf{Trunk}}(\square_n,-)\big)(\mathrm{inc})$ with $\mathrm{inc}\in\mathrm{Hom}_{\mathbf{Trunk}}\big(\mathrm{Conj(G)},\mathrm{Tr}(G_{\mathbf{Cat}})\big)$ is the inclusion of Conj(G) into $\mathrm{Tr}(G_{\mathbf{Cat}})$,
- $T^n = \mathrm{Hom}_{\mathbf{Set}}(-,A)\big(\theta(\square_n,G_{\mathbf{Cat}})\big)$ with $\theta(\square_n,G_{\mathbf{Cat}})$ the bijection induced by the adjunction $\mathrm{Cat}\overset{\theta}{\vdash}\mathrm{Tr}$ (cf. (4)),
- $\Sigma^n = \displaystyle\sum_{\sigma\in\mathbb{S}_n}\epsilon(\sigma)\,s_\sigma$ with $s_\sigma = \big(\mathrm{Hom}_{\mathbf{Set}}(-,A)\circ\mathrm{Hom}_{\mathbf{Cat}}(-,G_{\mathbf{Cat}})\big)(\sigma)$ where $\sigma\in\mathrm{Hom}_{\mathbf{Cat}}(\Delta_n,\square_n)$ is the functor defined in Example (3.2).

Theorem 5.6. *Let G be a group and A be an abelian group (considered as a trivial G-module). The graded linear map $S^\bullet$ is a chain complex morphism.*

Proof. The map $I^\bullet$ and $T^\bullet$ are induced by cubical set morphisms, hence they are chain complex morphisms. It remains to prove that $\Sigma^\bullet$ is a chain complex morphism.

We want to prove that $d_\square^{n+1}\big(\Sigma^n(f)\big) = \Sigma^{n+1}\big(d_\Delta^{n+1}(f)\big)$. The left term of this equation is equal to the sum of two terms (A) and (B) with

$$\text{(A)} = \sum_{i=1}^{n+1}\sum_{\sigma\in\mathbb{S}_n}(-1)^{i+1}\epsilon(\sigma)\,f\circ(\partial_{i,1}^{n+1}\circ\sigma)^*,$$
$$\text{(B)} = \sum_{i=1}^{n+1}\sum_{\sigma\in\mathbb{S}_n}(-1)^{i}\epsilon(\sigma)\,f\circ(\partial_{i,0}^{n+1}\circ\sigma)^*.$$

The right term is equal to sum of three terms (a), (b) and (c) with

$$\text{(a)} = \sum_{\sigma\in\mathbb{S}_{n+1}}\epsilon(\sigma)\,f\circ(\sigma\circ\partial_0^{n+1}),$$
$$\text{(b)} = \sum_{\sigma\in\mathbb{S}_{n+1}}(-1)^{n+1}\epsilon(\sigma)\,f\circ(\sigma\circ\partial_{n+1}^{n+1}),$$
$$\text{(c)} = \sum_{\sigma\in\mathbb{S}_{n+1}}\sum_{i=1}^{n}(-1)^{i}\epsilon(\sigma)\,f\circ(\sigma\circ\partial_i^{n+1}).$$

In order to prove the wanted equality we are going to show that

$$\text{(a)} = \text{(A)},\ \text{(b)} = \text{(B)}\ \text{and}\ \text{(c)} = 0.$$

•(a) = (A) : This equality is proved using the change of variables

$$\nu : \mathbb{S}_n\times\{1,\dots,n+1\}\longrightarrow\mathbb{S}_{n+1}$$

defined by $\nu(\sigma,i) := (i\,i-1\dots1)\circ(1\star\sigma)$. Using this change of variables, we have to prove for all $\sigma\in\mathbb{S}_n$ and $i\in\{1,\dots,n+1\}$:

1. $(\partial_{i,1}^{n+1}\circ\sigma)^* = (\nu(\sigma,i)\circ\partial_0^{n+1})^*$,
2. $(-1)^{i+1}\epsilon(\sigma) = \epsilon\big(\nu(\sigma,i)\big)$.

1. Let $\sigma \in \mathbb{S}_n$ and $i \in \{1, \dots, n+1\}$. We have $(\partial_{i,1}^{n+1} \circ \sigma)(k) = \sigma([k])_{<i} \amalg t_{+1}(\sigma([k])_{\geq i}) \amalg \{i\}$ and

$$\begin{aligned}
(\nu(\sigma, i) \circ \partial_0^{n+1})(k) &= \nu(\sigma, i)(k+1), \\
&= (i\, i-1 \dots 1)(\{1, 1+\sigma(1), \dots, 1+\sigma(k)\}, \\
&= \{i\} \amalg t_{-1}\big((1+\sigma([k]))_{<i}\big) \amalg t_{+1}(\sigma([k]))_{\geq i}), \\
&= \sigma([k])_{<i} \amalg t_{+1}(\sigma([k])_{\geq i}) \amalg \{i\}.
\end{aligned}$$

2. Clear.

•(b) = (B) : This equality is proved using the change of variables

$$\xi : \mathbb{S}_n \times \{1, \dots, n+1\} \longrightarrow \mathbb{S}_{n+1}$$

defined by $\xi(\sigma, i) := (i\, i+1 \dots n+1) \circ (\sigma \star 1)$. Using this change of variables, we have to prove for all $\sigma \in \mathbb{S}_n$ and $i \in \{1, \dots, n+1\}$:

1. $(\partial_{i,0}^{n+1} \circ \sigma)^* = (\xi(\sigma, i) \circ \partial_{n+1}^{n+1})^*$,
2. $(-1)^i \epsilon(\sigma) = (-1)^{n+1} \epsilon\big(\xi(\sigma, i)\big)$.

1. Let $\sigma \in \mathbb{S}_n$ and $i \in \{1, \dots, n+1\}$. We have $(\partial_{i,0}^{n+1} \circ \sigma)(k) = \sigma([k])_{<i} \amalg t_{+1}(\sigma([k])_{\geq i})$ and

$$\begin{aligned}
(\xi(\sigma, i) \circ \partial_{n+1}^{n+1})(k) &= \xi(\sigma, i)(k), \\
&= (i\, i+1 \dots n+1)(\{\sigma(1), \dots, \sigma(k)\}, \\
&= \sigma([k])_{<i} \amalg t_{+1}(\sigma([k]))_{\geq i}).
\end{aligned}$$

2. Clear.

•(c) = 0 : This equality is proved using the change of variables

$$\kappa : \{\sigma \in \mathbb{S}_{n+1} \,|\, \epsilon(\sigma) = 1\} \times \{1, \dots, n\} \longrightarrow \{\sigma \in \mathbb{S}_{n+1} \,|\, \epsilon(\sigma) = -1\} \times \{1 \dots, n\}$$

defined by $\kappa(\sigma, i) := \big((\sigma(i)\, \sigma(i+1)) \circ \sigma, i\big)$. Using this change of variables, we have to prove for all $\sigma \in \mathbb{S}_{n+1}$, $\epsilon(\sigma) = 1$ and $i \in \{1, \dots, n\}$:

1. $(\sigma \circ \partial_i^{n+1})^* = (\kappa(\sigma, i) \circ \partial_i^{n+1})^*$,
2. $(-1)^i \epsilon(\sigma) + (-1)^i \epsilon\big(\kappa(\sigma, i)\big)$.

1. Let $\sigma \in \mathbb{S}_{n+1}$ and $i \in \{1, \dots, n\}$. We have

$$(\kappa(\sigma, i) \circ \partial_i^{n+1})(k) = \begin{cases} \kappa(\sigma, i)(k) & \text{if } k < i, \\ \kappa(\sigma, i)(k+1) & \text{if } k \geq i. \end{cases} = \begin{cases} \sigma([k]) & \text{if } k < i, \\ \sigma([k+1]) & \text{if } k \geq i. \end{cases} = (\sigma \circ \partial_i^{n+1})(k).$$

2. Clear. □

Explicit formula for $S^\bullet$. By definition, for all $f \in C^n(G, A)$

$$S^n(f) = \sum_{\sigma \in \mathbb{S}_n} \epsilon(\sigma)\, f \circ \lambda \circ \sigma^* \circ \theta \circ \mathrm{inc}_* \circ \eta^{-1}.$$

Then to have an explict formula for $S^\bullet$ we have to compute

$$\lambda \circ \sigma^* \circ \theta \circ \mathrm{inc}_* \circ \eta^{-1} : \mathrm{Conj}(G)^n \to G^n$$

Let $(x_1,\dots,x_n)\in \mathrm{Conj}(G)^n$,

$$(\lambda\circ\sigma^*\circ\theta\circ \mathrm{inc}_*\circ\eta^{-1})(x_1,\dots,x_n)=(y_1,\dots,y_n),$$

where $y_k=(F\circ\sigma)(k-1\to k)$ for all $1\le k\le n$ with $F=(\eta^{-1})(x_1,\dots,x_n)$. Thus using Lemma (3.10) we find

$$\begin{aligned} y_k &= (F\circ\sigma)(k-1\to k),\\ &= F\big(\sigma([k-1])\to\sigma([k])\big),\\ &= \prod_{\substack{1\le x\le\sigma(k)\\ x\notin\sigma([k-1])}} F([x-1]\to[x]),\\ &= x_{i_1}\rhd\cdots\rhd x_{i_j}\rhd x_{\sigma(k)}, \end{aligned}$$

with $x_{i_1}<\cdots<x_{i_j}<\sigma(k)$ and $i_l\in\{\sigma(k+1),\dots,\sigma(n)\}$.

Theorem 5.7. *Let G be a group and A be an abelian group (considered as a trivial G-module). The graded abelian group morphism $S^\bullet$ from $C^\bullet(G,A)$ to $CR^\bullet(\mathrm{Conj}(G),A)$ defined by*

$$S^n(f)(x_1,\dots,x_n):=\sum_{\sigma\in\mathbb{S}_n}\epsilon(\sigma)\,f(y_1,\dots,y_n)$$

with $y_k=x_{i_1}\rhd\cdots\rhd x_{i_j}\rhd x_{\sigma(k)}$ where $x_{i_1}<\cdots<x_{i_j}<\sigma(k)$ and $i_l\in\{\sigma(k+1),\dots,\sigma(n)\}$, is a cochain complex.

$S^\bullet$ is a differential graded associative algebra morphism. Let G be a group and A be an associative algebra (considered as a trivial G-module). There is two differential graded associative algebras :

- $\{C^n(G,A),d^n_G,\cup\}_{n\in\mathbb{N}}$,
- $\{CR^n(\mathrm{Conj}(G),A),d^n_R,\star=\succ+\prec\}_{n\in\mathbb{N}}$.

and a cochain complex morphism $\{S^n\}_{n\in\mathbb{N}}$ between these two cochain complexes is defined in (16). The following theorem states that $\{S^n\}_{n\in\mathbb{N}}$ repects their associative structures.

Theorem 5.8. *Let G be a group and A be an associative algebra (considered as a trivial G-module). The cochain complex morphism*

$$\{S^n\}_{n\in\mathbb{N}}:\{C^n(G,A),d^n_G\}_{n\in\mathbb{N}}\longrightarrow\{CR^n(\mathrm{Conj}(G),A),d^n_R\}_{n\in\mathbb{N}}$$

is a differential graded associative algebra morphism. Hence it induces a graded associative algebra morphism $\{[S^n]\}_{n\in\mathbb{N}}$ from $\{H^n(G,A),\cup\}_{n\in\mathbb{N}}$ to $\{HR^n(\mathrm{Conj}(G),A),\star\}_{n\in\mathbb{N}}$.

Proof. By construction $\lambda^\bullet$ and $\eta^\bullet$ are graded associatve algebra morphism, hence we have to prove that $I^\bullet\circ T^\bullet\circ\Sigma^\bullet$ is a graded associative algebra morphism.

Let $f_1\in C^{p_1}_\Delta(G,A)$ and $f_2\in C^{p_2}_\Delta(G,A)$, on one hand

$$(I^{p_1+p_2}\circ T^{p_1+p_2}\circ\Sigma^{p_1+p_2})(f_1\cup_\Delta f_2):=\sum_{\sigma\in\mathbb{S}_{p_1+p_2}}\epsilon(\sigma)\,\mu_A\circ(f_1\times f_2)\circ\rho\circ\sigma^*\circ\theta\circ\mathrm{inc}_*,$$

and on the other hand

$$(I^{p_1}\circ T^{p_1}\circ \Sigma^{p_1})(f_1)\star(I^{p_2}\circ T^{p_2}\circ \Sigma^{p_2})(f_2) := \sum_{\gamma\in \mathrm{Sh}_{p_1,p_2}}\sum_{\alpha\in\mathbb{S}_{p_1}}\sum_{\beta\in\mathbb{S}_{p_2}} \epsilon(\gamma)\epsilon(\alpha)\epsilon(\beta)\,\mu_A\circ(f_1\times f_2)\circ\big((\alpha^*\circ\theta\circ \mathrm{inc}_*)\times(\beta^*\circ\theta\circ\mathrm{inc}_*)\big)\circ\rho_\sigma.$$

To show the equality of these two sums we are going to use the change of variables

$$\varphi : \mathrm{Sh}_{p_1,p_2}\times\mathbb{S}_{p_1}\times\mathbb{S}_{p_2}\longrightarrow \mathbb{S}_{p_1+p_2}$$

defined by $\varphi(\gamma,\alpha,\beta)=\gamma\circ(\alpha\star\beta)$. Using this change of variables we have to prove for all $\gamma\in\mathrm{Sh}_{p_1,p_2}$, $\alpha\in\mathbb{S}_{p_1}$ and $\beta\in\mathbb{S}_{p_2}$:

1. $\rho\circ\sigma^*\circ\theta\circ\mathrm{inc}_* = \big((\alpha^*\circ\theta\circ\mathrm{inc}_*)\times(\beta^*\circ\theta\circ\mathrm{inc}_*)\big)\circ\rho_\sigma$,
2. $\epsilon(\gamma)\epsilon(\alpha)\epsilon(\beta)=\epsilon\big(\gamma\circ(\alpha\star\beta)\big)$.

1. Let $\gamma\in\mathrm{Sh}_{p_1,p_2}$, $\alpha\in\mathbb{S}_{p_1}$ and $\beta\in\mathbb{S}_{p_2}$, we have to prove for all $F\in N\big(\mathrm{Conj}(G)\big)(\square_{p_1+p_2})$ the following equalities :

$$\begin{cases} \big(F\circ\sigma\circ(\alpha\star\beta)\circ j_{p_1}\big)(k-1\to k) = \big(F\circ\sigma\circ i_{p_1}\circ\alpha\big)(k-1\to k) & \text{if } 1\le k\le p_1,\\ \big(F\circ\sigma\circ(\alpha\star\beta)\circ j_{p_2}\big)(k-1\to k) = \big(F\circ\sigma\circ i_{p_2}\circ\beta\big)(k-1\to k) & \text{if } 1\le k\le p_2.\end{cases}$$

Let $1\le k\le p_1$,

$$\big(\sigma\circ(\alpha\star\beta)\circ j_{p_1}\big)(k) = (\sigma\circ(\alpha\star\beta)\big)(k) = (\sigma\circ(\alpha\star\beta)\big)([k]) = \sigma\big(\alpha([k])\big)$$

and

$$(\sigma\circ i_{p_1}\circ\alpha)(k) = (\sigma\circ i_{p_1})\big(\alpha([k])\big) = \sigma\big(\alpha([k])\big).$$

Let $1\le k\le p_2$,

$$\big(\sigma\circ(\alpha\star\beta)\circ j_{p_2}\big)(k) = (\sigma\circ(\alpha\star\beta)\big)(p_1+k) = (\sigma\circ(\alpha\star\beta)\big)([p_1+k]) = \sigma\big(t_{p_1}\beta([k])\big)$$

and

$$(\sigma\circ i_{p_2}\circ\beta)(k) = (\sigma\circ i_{p_2})\big(\beta([k])\big) = \sigma\big(t_{p_1}\beta([k])\big).$$

2. Clear. ☐

References

[Cov11] S. Covez. The local integration of Leibniz algebras. 2011. http://arxiv.org/abs/1011.4112.

[CS] J. S. Carter and M. Saito. Quandle homology theory and cocycle knot invariants. math.GT/0112026.

[Cuv] C. Cuvier. Algèbres de Leibniz: définitions, propriétés. *Ann. Sci. École Norm. Sup.*, 27 (1).

[EG03] P. Etingof and M. Graña. On rack cohomology. *Journal of pure and applied algebra*, 177:49–59, 2003.

[FR] R. Fenn and C. Rourke. Racks and links in codimension two. available at http://msp.warwick.ac.uk/ cpr/ftp/racks.ps.

[FRS95] R. Fenn, C. Rourke, and B. Sanderson. Trunks and classifying spaces. *Applied categorical structures*, 3:321–356, 1995.

[FT87] B. L. Feigin and B. L. Tsygan. Additive K-theory. *Lecture Notes in Math.*, 1289:67–209, 1987. in *K-theory, Arithmetic and Geometry (Moscow 1984-1986).*

[Kin07] M.K. Kinyon. Leibniz algebra, Lie racks and digroups. *J.Lie Theory*, 17:99–114, 2007.

[Lod93] J.-L. Loday. Une version non commutative des algèbres de Lie: les algèbres de Leibniz. *L'Enseignement Mathématique*, 39:269–293, 1993.

[Lod95] J-L. Loday. Cup-product for Leibniz cohomology and dual Leibniz algebra. *Math. Scand*, 77:189–196, 1995.

[Lod97] J.-L. Loday. Overview on Leibniz algebras, dialgebras and their homology. 17:91–102, 1997. in Cyclic cohomology and noncommutative geometry.

[Lod98] J.-L. Loday. *Cyclic Homology*, volume 301. Springer, 1998. second edition.

[Lod01] J.-L. Loday. Dialgebras. *Lecture Notes in Math.*, 1763:7–66, 2001. in Dialgebras and related operads.

[Lod03] J-L. Loday. Algebraic K-theory and the Conjectural Leibniz K-Theory. *K-Theory*, 30:105–127, 2003.

[Lod12] J.-L. Loday. Some problems in operad theory. *Proc. Int. Conf. in Nankai Series in Pure, Applied Mathematics and Theoretical Physics*, 9:139–146, 2012.

[LQ81] J.-L. Loday and D. Quillen. Cyclic homology and the Lie algebra homology of matrices. *Comment. Math. Helvet.*, 59 (4):569–591, 1981.

[Mac98] S. Maclane. *Categories for the working mathematician*, volume 5. Springer-Verlag, 1998. second edition.

[PS] J.H. Przytycki and A.S. Sikora. Distributive products and their homology. available at http://arxiv.org/abs/1105.3700.